



(19) Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada

Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

(11) CA 2 367 461 (13) A1

(40) 21.09.2000

(43) 21.09.2000

(12)

(21) 2 367 461

(22) 02.03.2000

(51) Int. Cl.<sup>7</sup>:

C07D 413/12, C07D 257/04,  
A61P 3/04, A61P 5/08,  
A61P 19/10, C07D 401/12,  
C07D 403/12, C07D 401/14,  
C07D 403/14, A61K 31/41,  
A61K 31/4178, A61K 31/4439,  
A61P 5/48, A61K 31/5377

(85) 10.09.2001

(86) PCT/US00/05704

(87) WO00/54729

(30) 60/124,131 US 12.03.1999  
60/154,919 US 21.09.1999

(71) BRISTOL-MYERS SQUIBB COMPANY,  
Lawrenceville-Princeton Road  
P.O. Box 4000, PRINCETON, XX (US).

LI, JUN (US).  
SWARTZ, STEPHEN G. (US).  
ROBL, JEFFREY (US).  
TINO, JOSEPH A. (US).  
HERNANDEZ, ANDRES S. (US).  
LI, JAMES J. (US).

(72)





(74) GOWLING LAFLEUR HENDERSON LLP

(54) COMPOSES HETEROCYCLIQUES AROMATIQUES UTILISES COMME SECRETAGOGUES D'HORMONES DE CROISSANCE

(54) HETEROCYCLIC AROMATIC COMPOUNDS USEFUL AS GROWTH HORMONE SECRETAGOGUES

(57)

Heterocyclic aromatic compounds are provided which are useful in stimulating endogenous production or release of growth hormone and in treating obesity, osteoporosis (improving bone density) and in improving muscle mass and muscle strength. The heterocyclic aromatic compounds have structure (I) including pharmaceutically acceptable salts thereof and all stereoisomers thereof, wherein X<sub>a</sub> is heteroaryl, preferably (a), (b) or (c) and R<sub>1</sub>, R<sub>1a</sub>, R<sub>6</sub>, Y, X<sub>b</sub>, A, B, Z, R<sub>3</sub>, R<sub>4</sub>, R<sub>4a</sub>, R<sub>5</sub> and R<sub>5a</sub> are as defined herein.





Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2367461 A1 2000/09/21

(21) 2 367 461

(12) DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2000/03/02  
(87) Date publication PCT/PCT Publication Date: 2000/09/21  
(85) Entrée phase nationale/National Entry: 2001/09/10  
(86) N° demande PCT/PCT Application No.: US 2000/005704  
(87) N° publication PCT/PCT Publication No.: 2000/054729  
(30) Priorités/Priorities: 1999/03/12 (60/124,131) US;  
1999/09/21 (60/154,919) US

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C07D 413/12, A61K 31/5377, A61P 5/48,  
A61P 5/08, A61P 3/04, A61K 31/4439, A61K 31/4178,  
A61K 31/41, A61P 19/10, C07D 401/12, C07D 401/14,  
C07D 403/12, C07D 403/14, C07D 257/04





(71) Demandeur/Applicant:  
BRISTOL-MYERS SQUIBB COMPANY, US

(72) Inventeurs/Inventors:  
SWARTZ, STEPHEN G., US;  
LI, JUN, US;  
LI, JAMES J., US;  
HERNANDEZ, ANDRES S., US;

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSES HETEROCYCLIQUES AROMATIQUES UTILISES COMME SECRETAGOGUES D'HORMONES  
DE CROISSANCE

(54) Title: HETEROCYCLIC AROMATIC COMPOUNDS USEFUL AS GROWTH HORMONE SECRETAGOGUES



(57) Abrégé/Abstract:

Heterocyclic aromatic compounds are provided which are useful in stimulating endogenous production or release of growth hormone and in treating obesity, osteoporosis (improving bone density) and in improving muscle mass and muscle strength. The heterocyclic aromatic compounds have structure (I) including pharmaceutically acceptable salts thereof and all stereoisomers thereof, wherein X<sub>a</sub> is heteroaryl, preferably (a), (b) or (c) and R<sub>1</sub>, R<sub>1a</sub>, R<sub>6</sub>, Y, X<sub>b</sub>, A, B, Z, R<sub>3</sub>, R<sub>4</sub>, R<sub>4a</sub>, R<sub>5</sub> and R<sub>5a</sub> are as defined herein.

Canada

<http://opic.gc.ca> · Ottawa-Hull K1A 0C9 · <http://cipo.gc.ca>

OPIC · CIPO 191

OPIC



CIPO

(72) Inventeurs(suite)/Inventors(continued): TINO, JOSEPH A., US; ROBL, JEFFREY, US

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
21 September 2000 (21.09.2000)

PCT

(10) International Publication Number  
WO 00/54729 A3(51) International Patent Classification<sup>7</sup>: A61K 31/41,  
31/4178, 31/4439, 31/5377, A61P 3/04, 5/08, 5/48, 19/10,  
C07D 401/12, 401/14, 403/12, 403/14, 413/12, 413/14(74) Agents: RODNEY, Burton et al.; Bristol-Myers Squibb  
Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(21) International Application Number: PCT/US00/05704

(22) International Filing Date: 2 March 2000 (02.03.2000)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:  
60/124,131 12 March 1999 (12.03.1999) US  
60/154,919 21 September 1999 (21.09.1999) US(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY  
[US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).(72) Inventors: ROBL, Jeffrey; 7 Tulip Drive, Newtown,  
PA 18940 (US). TINO, Joseph, A.; 11 Chopin Lane,  
Lawrenceville, NJ 08648 (US). HERNANDEZ, Andres,  
S.; 1102 White Pine Circle, Lawrenceville, NJ 08648  
(US). LI, James, J.; 76 Stanford Road East, Pennington,  
NJ 08534 (US). LI, Jun; 137 York Drive, Princeton, NJ  
08540 (US). SWARTZ, Stephen, G.; 162 Willow Lane,  
Warrington, PA 18976 (US).(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA,  
UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

With international search report.

(88) Date of publication of the international search report:  
11 January 2001For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLIC AROMATIC COMPOUNDS USEFUL AS GROWTH HORMONE SECRETAGOGUES

(57) Abstract: Heterocyclic aromatic  
compounds are provided which are  
useful in stimulating endogenous  
production or release of growth hormone  
and in treating obesity, osteoporosis  
(improving bone density) and in  
improving muscle mass and muscle  
strength. The heterocyclic aromatic  
compounds have structure (I) including  
pharmaceutically acceptable salts thereof  
and all stereoisomers thereof, wherein  
 $X_a$  is heteroaryl, preferably (a), (b) or (c)  
and  $R_1$ ,  $R_{1a}$ ,  $R_6$ ,  $Y$ ,  $X_b$ ,  $A$ ,  $B$ ,  $Z$ ,  $R_3$ ,  $R_4$ ,  
 $R_{4a}$ ,  $R_5$  and  $R_{6a}$  are as defined herein.




Summary of the Invention

In accordance with the present invention, novel heterocyclic aromatic compounds are provided which are growth hormone secretagogues and have the structure

5

I




including pharmaceutically acceptable salts thereof, prodrug esters thereof, and all stereoisomers thereof, wherein  $R_1$  is alkyl, aryl, alkenyl, alkynyl,


- 10 arylalkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, arylalkyloxyalkyl, aryloxyalkyl, heteroaryl, cycloalkylalkoxyalkyl, heteroarylalkyl, cycloheteroalkyl or cycloheteroalkylalkyl (where the above groups are defined below) and these groups may be optionally substituted by 1,2 or 3-substituents selected from
- 15 halogen,  $-OR_8$ ,  $-OC(O)R_8$ , alkyl, phenyl, phenoxy, halophenyl,  $-CF_3$ ,  $-OCF_3$ ,  $-N(R_{8a})C(O)(R_8)$ , or  $-N(R_8)(R_{8a})$ ;

$R_{1a}$  is H, alkyl, or cycloalkyl;

$X_a$  is heteroaryl, which preferably include

20





A is oxygen, sulfur, -NH-, -N-R<sub>5</sub>, or -NC(O)-R<sub>2</sub>;

B is -CR<sub>5b</sub> or -N-;

Z is a bond or -S-;

5 G is oxygen or sulfur;

U is oxygen, sulfur, -NH-, or -N-R<sub>5b</sub>;

R<sub>2</sub> is alkyl, aryl, alkenyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, heteroaryl, or heteroarylalkyl (where the above groups are  
10 defined below) and these groups may optionally be substituted by 1,2 or 3-substituents selected from halogen, -OR<sub>8b</sub>, -OC(O)R<sub>8b</sub>, alkyl, phenyl, phenoxy, halophenyl, -CF<sub>3</sub>, -OCF<sub>3</sub>, -N(R<sub>8c</sub>)C(O)(R<sub>8b</sub>), or -N(R<sub>8c</sub>)(R<sub>8b</sub>);

15 R<sub>3</sub> is H, halogen, alkyl, aryl, alkenyl, alkynyl, alkaryl, alkoxy, aryloxy or J1, and where alkyl, aryl, alkenyl, alkynyl, arylalkyl, alkoxy, or aryloxy may be optionally substituted with 1 to 3 J1;

R<sub>4</sub> and R<sub>4a</sub> are the same or different and are  
20 independently H, halogen, -CF<sub>3</sub>, alkyl, or aryl;


R<sub>5</sub> is H, alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, cycloalkylalkoxyalkyl, -SO<sub>2</sub>T<sub>1</sub>, -SO<sub>2</sub>N(T<sub>1a</sub>)T<sub>1</sub>, or  
25 heteroarylalkyl, and where alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, heteroaryloxyalkyl, cycloalkylalkoxyalkyl, or heteroarylalkyl may be  
30 independently optionally substituted with 1 to 3 J1;

R<sub>5a</sub> and R<sub>5b</sub> are the same or different and are independently H, alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl,

cycloalkylalkoxyalkyl, heteroarylalkyl, or J1, and where alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, heteroaryloxyalkyl,

- 5 cycloalkylalkoxyalkyl, or heteroarylalkyl may be independently optionally substituted with 1 to 3 J1;

Y is



- 10 where x and y are independently 0 to 3 and z is 1 to 3;

X<sub>c</sub> is a bond, -N-R<sub>6a</sub> or -O-;


R<sub>7</sub> and R<sub>7a</sub> are the same or different and are independently H, alkyl, -CF<sub>3</sub>, phenyl, aryl, arylalkyl, and cycloalkyl; or one or both of R<sub>7</sub> and R<sub>7a</sub> can be

- 15 independently joined to one or both of R<sub>9</sub> and R<sub>10</sub> groups (of X<sub>b</sub>) to form an alkylene bridge of 1 to 5 carbon atoms; or R<sub>7</sub> and R<sub>7a</sub> are joined together to form a ring of from 3-7 carbon atoms;

- 20 R<sub>6</sub>, R<sub>6a</sub>, R<sub>6b</sub>, R<sub>6c</sub>, R<sub>8</sub>, R<sub>8a</sub>, R<sub>8b</sub>, R<sub>8c</sub>, R<sub>8d</sub>, R<sub>8e</sub>, R<sub>8f</sub>, R<sub>8g</sub>, R<sub>8h</sub>, R<sub>8i</sub>, R<sub>8k</sub>, R<sub>8l</sub>, and R<sub>8m</sub> are the same or different and are independently H, alkyl, cycloalkyl, alkenyl or aryl;

R<sub>8j</sub> is H, alkyl, aryl, hydroxy or -OC(O)R<sub>8k</sub>;

X<sub>b</sub> is



25

R<sub>9</sub> and R<sub>10</sub> are the same or different and are independently selected from H, alkyl, and substituted alkyl where the substituents may be 1 to 3 hydroxys, 1 to 3 C<sub>1</sub>-C<sub>10</sub>-alkanoyloxy; 1 to 3 C<sub>1</sub>-6 alkoxy, phenyl, phenoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl; or R<sub>9</sub> and R<sub>10</sub> can together form

- 30



-(CH<sub>2</sub>)<sub>t</sub>X<sub>d</sub>(CH<sub>2</sub>)<sub>u</sub>- where X<sub>d</sub> is C(R<sub>8h</sub>)(R<sub>8j</sub>), -O- or -N(R<sub>6b</sub>),  
t and u are independently 1-3;

R<sub>11</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, -CF<sub>3</sub>, arylalkyl, or aryl, and  
with the alkyl and aryl groups being optionally


5 substituted with 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-<sub>10</sub>alkanoyloxy,  
1 to 3 C<sub>1</sub>-<sub>6</sub>alkoxy, phenyl, phenoxy or C<sub>1</sub>-<sub>6</sub>alkoxycarbonyl;

R<sub>12</sub> and R<sub>13</sub> are independently H, C<sub>1</sub>-C<sub>6</sub>alkyl, -CF<sub>3</sub>,  
aryl, or halogen, and with the alkyl and aryl groups being  
optionally substituted with 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-<sub>10</sub>-  
10 alkanoyloxy, 1 to 3 C<sub>1</sub>-<sub>6</sub>alkoxy, or C<sub>1</sub>-<sub>6</sub>alkoxycarbonyl;

J<sub>1</sub> is nitro, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>CN,  
-(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)N(T<sub>1b</sub>)T<sub>1</sub>,  
-(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)SO<sub>2</sub>T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)OT<sub>1</sub>,  
-(CH<sub>2</sub>)<sub>v</sub>OC(O)OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OC(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OC(O)N(T<sub>1a</sub>)T<sub>1</sub>,  
15 -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)SO<sub>2</sub>N(T<sub>1b</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>SO<sub>2</sub>T<sub>1</sub>,  
-(CH<sub>2</sub>)<sub>v</sub>SO<sub>2</sub>N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>CH(OH)T<sub>1</sub>,  
cycloheteroalkyl, or heteroaryl as defined below, with v  
being 0-5;

T<sub>1</sub>, T<sub>1a</sub> and T<sub>1b</sub> are the same or different and are  
20 independently H, alkyl, alkenyl, alkynyl, lower  
alkythioalkyl, alkoxyalkyl, aryl, arylalkyl, heteroaryl,  
heteroarylalkyl, cycloheteroalkyl, or cycloalkyl, each of  
which may be optionally substituted with 1, 2 or 3 of the  
following groups, halogen, hydroxyl, -NR<sub>8f</sub>C(O)NR<sub>8g</sub>R<sub>8i</sub>,  
25 -C(O)NR<sub>8f</sub>R<sub>8g</sub>, -NR<sub>8f</sub>C(O)R<sub>8g</sub>, -CN, -N(R<sub>8f</sub>)SO<sub>2</sub>R<sub>14</sub>, -OC(O)R<sub>8f</sub>,  
-SO<sub>2</sub>NR<sub>8f</sub>R<sub>8g</sub>, -SOR<sub>14</sub>, -SO<sub>2</sub>R<sub>14</sub>, alkoxy, -COOH,  
cycloheteroalkyl, or -C(O)OR<sub>14</sub>; or T<sub>1</sub> and T<sub>1a</sub> or T<sub>1</sub> and  
T<sub>1b</sub> can together form -(CH<sub>2</sub>)<sub>w</sub>X<sub>e</sub>(CH<sub>2</sub>)<sub>z</sub>- where X<sub>e</sub> is  
-C(R<sub>8m</sub>)(R<sub>8l</sub>), -O-, -S-, -SO-, -SO<sub>2</sub>-, -NC(O)OR<sub>14a</sub>,  
30 -NC(O)NR<sub>14a</sub>R<sub>14b</sub>, -NC(O)R<sub>14a</sub> or -N(R<sub>6c</sub>) where w and z are  
each independently 1-3; with the proviso that T<sub>1</sub> can not  
be hydrogen when it is connected to carbonyl or sulfur, as  
in -C(O)T<sub>1</sub> or -SO<sub>2</sub>T<sub>1</sub>;

R<sub>14</sub>, R<sub>14a</sub>, and R<sub>14b</sub> are independently C<sub>1</sub>-C<sub>6</sub>alkyl,  
35 heteroaryl, or aryl, each optionally substituted with  
-(CH<sub>2</sub>)<sub>s</sub>OH, with s being 0-5;




with the proviso that where  $X_a$  is

(1) where one or both of  $R_7$  and  $R_{7a}$ , and one or both of  $R_9$  and  $R_{10}$  form an alkylene bridge, then where  $R_5$  is

5  $-(CH_2)C(O)N(T_{1a})T_1$ , then at least one of  $T_{1a}$  and  $T$   $\neq$  H; or

(2) where  $R_1$  is arylalkyl and  $R_{1a}$  is H and  $R_5$  is  $-(CH_2)C(O)N(T_{1a})T_1$ , then  $T_{1a}$  or  $T_1$  is other than



10 (3) where  $R_1$  and  $R_7$  are each 2-naphthyl- $CH_2$ -, then  $R_5 \neq$  phenethyl.

The compounds of the instant invention all have at least one asymmetric center as noted by the asterisk in structural formula I. Additional asymmetric centers may be present on the molecule depending upon the nature of the various substituents on the molecule. Each such asymmetric center will produce two optical isomers and it is intended that all such optical isomers, as separated, pure or partially purified optical isomers or racemic mixtures thereof, be included within the ambit of the instant invention. In the case of the asymmetric center represented by the asterisk in formula I, the more active and thus more preferred configuration is R as determined by the R/S rules when  $R_{1a}$  is H. Isomers may be separated by conventional methods, for example, chromatographic or fractional crystallization.

The pharmaceutically acceptable salts of the compounds of formulae I of the invention include alkali metal salts such as lithium, sodium or potassium, alkaline earth metal salts such as calcium or magnesium, as well as zinc or aluminum and other cations such as ammonium,

choline, diethanolamine, ethylenediamine, t-butylamine, t-octylamine, dehydroabietylamine, as well as pharmaceutically acceptable anions such as phosphate, mandelate, chloride, bromide, iodide, tartrate, acetate, 5 methanesulfonate, maleate, succinate, glutarate, and salts of naturally occurring amino acids such as arginine, lysine, alanine and the like, and prodrug esters thereof.

In addition, in accordance with the present invention, a method for increasing levels of endogenous 10 growth hormone or increasing the endogenous production or release of growth hormone is provided wherein a compound of formula I as defined hereinbefore is administered in a therapeutically effective amount.

Furthermore, in accordance with the present 15 invention, a method is provided for preventing or treating osteoporosis (improving bone density and/or strength), or treating obesity, or increasing muscle mass and/or muscle strength, or maintenance of muscle strength and function in elderly humans, or reversal or prevention of frailty 20 in elderly humans, wherein a compound of formula I as defined hereinbefore is administered in a therapeutically effective amount.

Detailed Description of the Invention

The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

- 5 Unless otherwise indicated, the term "lower alkyl", "alkyl" or "alk" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons, containing 1 to 40 carbons, preferably 1 to 20 carbons, more preferably 1 to 6 carbons, in the normal
- 10 chain, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, the various branched chain isomers thereof, and the like as well as such groups
- 15 including 1 to 3 substituents including alkyl, aryl, alkenyl, alkynyl, hydroxy, arylalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, arylalkoxy, alkanoyl, amino, haloaryl, CF<sub>3</sub>, OCF<sub>3</sub>, aryloxy, heteroaryl, cycloalkylalkoxyalkyl, or cycloheteroalkyl.
- 20 Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and
- 25 tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 4 to 7 carbons, forming the ring and which may be fused to 1 aromatic ring as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclododecyl, cyclohexenyl,
- 30



any of which groups may be optionally substituted with 1 to 3 substituents as defined above for alkyl.

The term "aryl" as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl) and may optionally include one to three additional rings fused to "aryl" (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings) and may be optionally substituted through available carbon atoms with 1, 2, or 3 groups selected from hydrogen, halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkylalkyl, fluorenyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, arylthio, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, oxo, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfonaminocarbonyl, or preferably any of the aryl substituents as set out above.

Preferred aryl groups include phenyl, biphenyl or naphthyl.

The term "aralkyl", "aryl-alkyl" or "aryllower alkyl" as used herein alone or as part of another group refers to alkyl groups as discussed above having an aryl substituent, such as benzyl or phenethyl, or naphthylpropyl, or an aryl as defined above.

The term "lower alkoxy", "alkoxy", "aryloxy" or "aralkoxy" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to an oxygen atom.

The term "amino" as employed herein alone or as part of another group may optionally be substituted with one or two substituents such as alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and/or cycloalkyl.

The term "lower alkylthio", "alkylthio", "arylthio" or "aralkylthio" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to a sulfur atom.

The term "lower alkylamino", "alkylamino", "arylamino", or "arylalkylamino" as employed herein alone or as part of another group includes any of the above alkyl, aryl or arylalkyl groups linked to a nitrogen atom.

The term "acyl" as employed herein by itself or part of another group, as defined herein, refers to an organic radical linked to a carbonyl  $\left( \begin{smallmatrix} \text{O} \\ \parallel \\ \text{C} \end{smallmatrix} \right)$  group; examples of acyl groups include alkanoyl, alkenoyl, aroyl, aralkanoyl, heteroaroyl, cycloalkanoyl, and the like.

The term "alkanoyl" as used herein alone or as part of another group refers to alkyl linked to a carbonyl group.

Unless otherwise indicated, the term "lower alkenyl" or "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 3 to 12 carbons, and more preferably 2 to 6 carbons in the normal chain, which include one to six double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, hydroxy, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol,

alkylthio or any of the substituents for alkyl as set out herein.




Unless otherwise indicated, the term "lower alkynyl" or "alkynyl" as used herein by itself or as part  
5 of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one triple bond in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl,  
10 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl,  
15 amino, heteroaryl, cycloheteroalkyl, hydroxy, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, and/or alkylthio, or any of the substituents for alkyl as set out herein.

The term "alkylene" as employed herein alone or as  
20 part of another group refers to alkyl groups as defined above having single bonds for attachment to other groups at two different carbon atoms and may optionally be substituted as defined above for "alkyl".



The terms "alkenylene" and "alkynylene" as employed  
25 herein alone or as part of another group refer to alkenyl groups as defined above and alkynyl groups as defined above, respectively, having single bonds for attachment at two different carbon atoms.

Examples of  $(CH_2)_x$ ,  $(CH_2)_y$ ,  $(CH_2)_w$ ,  $(CH_2)_v$ ,  $(CH_2)_s$ ,  
30  $(CH_2)_t$ ,  $(CH_2)_u$  or  $(CH_2)_z$  groups (which may include alkylene, alkenylene or alkynylene groups as defined herein, and may optionally include 1, 2, or 3 substituents which may be any of the alkyl substituents set out herein), are as follows:

35



5



10



15           The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine as well as CF<sub>3</sub>, with chlorine or fluorine being preferred.


20           The term "metal ion" refers to alkali metal ions such as sodium, potassium or lithium and alkaline earth metal ions such as magnesium and calcium, as well as zinc and aluminum.

25           The term "heterocyclic", "heterocyclo" or "heterocycle" as employed herein alone or as part of another group refers to "heteroaryl" groups or "cycloheteroalkyl" groups.

          The term "cycloheteroalkyl" as used herein alone or as part of another group refers to a 4-, 5-, 6- or 7-





membered saturated or partially unsaturated ring which includes 1 to 2 hetero atoms such as nitrogen, oxygen and/or sulfur, linked through a carbon atom or a heteroatom, where possible, optionally via the linker  
 5 (CH<sub>2</sub>)<sub>p</sub> (which is defined above), such as



and the like. The above groups may include 1 to 4 substituents such as alkyl, halo, oxo and/or any of of the  
 15 aryl substituents set out herein. In addition, any of the above rings can be fused to a cycloalkyl, aryl, heteroaryl or cycloheteroalkyl ring.

The term "heteroaryl" or "heterocyclicaryl" as used herein alone or as part of another group refers to a 5- or  
 20 6- membered aromatic ring which includes 1, 2, 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur, and such rings fused to an aryl, cycloalkyl, heteroaryl or cycloheteroalkyl ring (e.g. benzothiophenyl, indolyl), and includes possible N-oxides, such as





5

and the like.

The heteroaryl groups may optionally include 1 to 4 substituents such as any of the aryl substituents set out herein. In addition, any of the above rings can be fused  
 10 to a cycloalkyl, aryl, heteroaryl or cycloheteroalkyl ring.

The term "prodrug esters" of the formula I compounds includes esters of hydroxyls and phenols, such as acetate, benzoate, pivalate, stearoylate, isobutyrate,  
 15 and the like as known in the art.

#### General Synthetic Schemes

The compounds of the present invention may be prepared according to the following general synthetic  
 20 schemes, as well as relevant published literature procedures that are used by the one skilled in the art. Exemplary reagents and procedures for these reactions appear hereinafter and in the working Examples. Unless otherwise specified, the various substituents of the  
 25 compounds are defined in the same manner as the formula I compound of the invention.

In the following reactions, amide bond forming (peptide coupling) reactions are conducted under standard peptide coupling procedures known in the art. Optimally,  
 30 the reaction is conducted in a solvent such as dimethylformamide (DMF) at 0°C to room temperature using

1-ethyl-3-(3-dimethylamino-propyl)carbodiimide (EDAC) or (WSC)), 1-hydroxybenzo-triazole hydrate (HOBt) or 1-hydroxy-7-azabenzotriazole (HOAt) and a base, for example 4-dimethylaminopyridine (DMAP) or Hunig's base.

- 5           Unnatural protected amino acids can be purchased or prepared by standard methods known in the art. Chiral preparations include Meyers, A. G. et al., *J. Am. Chem. Soc.*, 119, 656-673 (1997). O-Alkylated serine derivatives can be formed from serine by known methods, Maligres, P. E. et al., *Tetrahedron*, 53, 10983-10992 (1997).

- The cyclizations to benzimidazoles or benzoxazoles in Schemes 1, 1a, and 1b can be carried out under standard conditions known in the art. Suitable cyclization procedures are described in Nestor, Jr., J.J., *J. Med. Chem.*, 27, 320-5 (1984) and are optimally heating (60-90°C) in an acidic solvent, such as acetic acid. Benzothiazoles can be prepared as in Spitulnik, M. J., *Synthesis*, 730 (1976), and by other methods known in the art.

- 20           Protection and deprotection in the Schemes below may be carried out by procedures generally known in the art. See, for example, T. W. Greene, *Protecting Groups in Organic Synthesis*, Second Edition, 1991. PG in the Schemes below denotes a nitrogen protecting group, optimally BOC.
- 25           The BOC group can be removed under acidic conditions, optimally HCl or trifluoroacetic acid.


- Reduction of nitro groups in Schemes 1c and 1d can be carried out by methods known in the art. Optimally the reduction can be carried out with H<sub>2</sub> in the presence of a catalyst (palladium or platinum).

- 30           The arylation reaction covered in Scheme 2 can be performed under the known Suzuki coupling conditions in the literature, or other conditions using the metals zinc or tin known in the art. If the Rx is a chloro, the coupling can be performed as disclosed in Indolese, A. F., *Tet. Lett.*, 38, 3513-3516 (1997) or Shen, W. *Tet. Lett.*, 38, 5575-5578 (1997) using a nickel catalyst.

5       The triazole forming reaction found in Scheme 4 can  
be carried out under standard conditions known in the art.  
Suitable procedures are described in Sung, K et al.  
*Heterocyc. Chem.*, 29, 1101 (1992), Shukla, J.S. et al.  
*Indian J Chem*,, 30B, 332 (1991), and Wilson, M. W. et al.  
10 *Molecular Diversity*, 3, 95 (1998).

The transformation of alcohols 69 to azido 70 and 75 to 76 (Schemes 5 and 5a) can be performed by methods known in the art. A single pot procedure described in Thompson, A.S. et al. *J Org Chem.*, 58, 886 (1993), is the optimal transformation.

The tetrazole forming reaction shown in Scheme 6 can be carried out on solid phase resin.

**polystyrene**-


The resin may be a Merrifield type resin.

Intermediates 28 and 42 can be further transformed as shown previously in Schemes 3. Alternatively, intermediates 28 and 42 can be treated with an acid linked  
5 to a resin, such as Merrifield or Rink, to give resin bound amides, which can be cleaved to give compounds VI and IX. The carbamate resin linker depicted in Scheme 6 is only one possible candidate (Hernandez, A. et al. *J. Org. Chem.*, 62, 3153 (1997)).


10 Compounds in all Schemes except 1b with a terminal -NH-R<sub>9</sub> moiety can be transformed by methods known in the art, such as reductive amination or alkylation, to

compounds of the form  $\text{—}\overset{\text{R}_9}{\underset{\text{R}_{10}}{\text{N}}}$ .


Scheme 1



Scheme 1a






Scheme 1b






Scheme 1c




$Y_a = \text{O}, \text{S}, \text{NH}$

$Z = -\text{CO}, -\text{SO}_2$




Scheme 1d




Scheme 1e




Scheme 2








## Scheme 3a: Tetrazole




## Scheme 3b



(where  $R_5$  is a protecting group PGx)





(where  $R_5$  is a protecting group)




where  $R_9$  and  $R_{10}$  are other than hydrogen

or




[illegible]

IX






## Scheme 3d



29



Where  $R_5 = -CH_2(CH_2)_nCO_2R_x$   
 $n = 0-4$

43




29a


Where  $R_5 = -CH_2(CH_2)_nCO_2R_x$   
 $n = 0-4$



## Scheme 3e




Where  $R_5 = -CH_2(CH_2)_n-N(T_{1a})-PG_x$   
 $n = 0-4$ ,  $PG_x$  = a protecting group




Where  $R_5 = -CH_2(CH_2)_n-N(T_{1a})-PG_x$   
 $n = 0-4$ ,  $PG_x$  = a protecting group

5

## Scheme 3f




Where  $R_5 = -CH_2(CH_2)_n-N(T_{1a})-PG_x$   
 $n = 0-4$ ,  $PG_x$  = a protecting group




Where  $R_5 = -CH_2(CH_2)_n-N(T_{1a})-PG_x$   
 $n = 0-4$ ,  $PG_x$  = a protecting group

## Scheme 3g




Where  $\text{R}_5 = -\text{CH}_2(\text{CH}_2)_n\text{O}-\text{PG}_x$   
 $n = 0-4$ ,  $\text{PG}_x =$  a protecting group




Where  $\text{R}_5 = -\text{CH}_2(\text{CH}_2)_n\text{O}-\text{PG}_x$   
 $n = 0-4$ ,  $\text{PG}_x =$  a protecting group

5

## Scheme 3h




Where  $\text{R}_5 = -\text{CH}_2(\text{CH}_2)_n\text{O}-\text{PG}_x$   
 $n = 0-4$ ,  $\text{PG}_x =$  a protecting group





Where  $\text{R}_5 = -\text{CH}_2(\text{CH}_2)_n\text{O}-\text{PG}_x$   
 $n = 0-4$ ,  $\text{PG}_x =$  a protecting group




## Scheme 4: Triazoles






## Scheme 4a




## 5 Scheme 4b




## Scheme 4c






## Scheme 4d



## Scheme 5: Imidazoles




$R_4$  and  $R_{4a}$  are independently H, halogen, alkyl, aryl, alkoxy, aryloxyalkyl, or J1




$R_{5b1}$  and  $R_{5b2}$  are independently alkyl or aryl






Scheme 5a



Scheme 6:



Alternatively, when  $R_5$  is  $-\text{CH}_2(\text{CH}_2)_n\text{CO}_2\text{R}_x$




The conditions described here for carrying out each step in the general synthetic schemes are conventional and capable of wide variation. They are presented for illustrative purpose only and are not intended as a restriction on the scope of invention.


Final compounds can be easily purified by recrystallization, silica gel chromatography, or reverse phase prep HPLC. In the cases where reverse phase prep HPLC is used, a mixture of solvent A (10% MeOH/90% H<sub>2</sub>O/0.2% TFA) and solvent B (90% MeOH/10 %H<sub>2</sub>O/0.2% TFA) are used.

Preferred compounds of formula I of the invention include compounds of the structure wherein Xa is as indicated:

IA




IB





5

IC




Preferred are compounds of the present invention of the structure I wherein R<sub>1</sub> is arylalkyl, arylalkyloxyalkyl, aryloxyalkyl, cycloheteroalkylalkyl, heteroarylalkyl, for example




R<sub>1a</sub> is H or alkyl;

15


and (1) X<sub>a</sub> is



where  $R_5$  is alkyl, alkenyl or heteroaryloxy-alkyl, each substituted with  $J_1$ , and  $J_1$  is  $-(CH_2)_vOC(O)N(T_{1a})T_1$ ,  $-(CH_2)_vCN$ , or heteroaryl; or




(2)  $X_a$  is where  $A$  is  $NH$ ,  $R_4$  and  $R_{4a}$  are  $H$  and  $R_3$  is  $J_1$ ; or




(3)  $X_a$  is where  $B$  is  $-N-$ ;  $Z$  is a bond or  $-S-$ ;  $R_{5a}$  is  $H$ , or alkyl or arylalkyl each substituted with 1 to 3  $J_1$ ;  $R_5$  is alkyl optionally substituted with  $J_1$ ;

10  $R_6$  is  $H$ ;



$Y$  is where  $x$  and  $y$  are 0,  $X_c$  is a bond, and  $R_7$  and  $R_{7a}$  are independently alkyl;



$X_b$  is

$R_{8j}$  is hydroxy or  $-OC(O)R_{8k}$  where  $R_{8k}$  is alkyl or

15 aryl;

$R_9$  and  $R_{10}$  are independently  $H$  and substituted alkyl where the substituents may be 1 or 2 hydroxyls;

$J_1$  is  $-(CH_2)_vCN$ ,  $-(CH_2)_vN(T_{1a})SO_2T_1$ ,


$-(CH_2)_vC(O)N(T_{1a})T_1$ ,  $-(CH_2)_vN(T_{1a})C(O)T_1$ ,

20  $-(CH_2)_vOC(O)N(T_{1a})T_1$ ,  $-(CH_2)_vN(T_{1a})C(O)N(T_{1b})T_1$ , or heteroaryl, with  $v$  being 0-4;

$T_1$ ,  $T_{1a}$  and  $T_{1b}$  are independently alkyl, lower alkythioalkyl, alkoxyalkyl, aryl, arylalkyl,


heteroarylalkyl, cycloheteroalkyl, or cycloalkyl, each of which may be optionally substituted with  $-\text{OC}(\text{O})\text{R}_{8f}$ ,  $-\text{C}(\text{O})\text{NR}_8\text{R}_{8a}$ ,  $-(\text{CH}_2)_s\text{OH}$ , with  $s$  being 0-2,  $-\text{SO}_2\text{NR}_8\text{R}_{8a}$ ,  $-\text{SO}_2\text{R}_{14}$ , or alkoxy; or  $\text{T}_1$  and  $\text{T}_{1a}$  or  $\text{T}_1$  and  $\text{T}_{1b}$  can  
 5 together form  $-(\text{CH}_2)_w\text{X}_e(\text{CH}_2)_z-$  where  $\text{X}_e$  is  $\text{C}(\text{R}_{8m})(\text{R}_{8l})$ ;  
 $\text{R}_{14}$  is  $\text{C}_1$ - $\text{C}_6$ alkyl optionally substituted with  $-(\text{CH}_2)_v\text{OH}$ , with  $v$  being 0-2;


Preferred compounds of the invention include the  
 10 following:






Chiral










5 More preferred compounds of the invention include the following:





10

The present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, at least one of the compounds of formula I in association with a pharmaceutical carrier or diluent.

5 Optionally, the active ingredient of the pharmaceutical compositions can comprise a growth promoting agent in addition to at least one of the compounds of formula I or another composition which exhibits a different activity, e.g., an antibiotic or other pharmaceutically active  
10 material.

Growth promoting agents include, but are not limited to, TRH, diethylstilbesterol, theophylline, enkephalins, E series prostaglandins, compounds disclosed in U.S. Patent No. 3,239,345, e.g., zeranol, and compounds  
15 disclosed in U.S. Patent No. 4,036,979, e.g., sulbenox or peptides disclosed in U.S. Patent No. 4,411,890.

A still further use of the disclosed compounds of formula I of the invention is in combination with other growth hormone secretagogues such as GHRP-6, GHRP-1 as  
20 described in U.S. Patent No. 4,411,890; and publications WO 89/07110 and WO 89/07111 and B-HT920 or growth hormone releasing factor and its analogs or growth hormone and its analogs or somatomedins including IGF-1 and IGF-2. A  
25 still further use of the disclosed compounds of formula I of the invention is in combination with parathyroid hormone or a bisphosphonate, such as MK-217 (alendronate), in the treatment of osteoporosis.

A still further use of the disclosed compounds of formula I is in combination with estrogen, testosterone, a  
30 selective estrogen receptor modulator, such as tamoxifen or raloxifene, or a selective androgen receptor modulator, such as disclosed in Edwards, J. P. et al., *Bio. Med. Chem. Let.*, 9, 1003-1008 (1999) and Hamann, L. G. et al., *J. Med. Chem.*, 42, 210-212 (1999), for the treatment of  
35 aspects of Metabolic Syndrome, maintenance of muscle strength and function in elderly humans, reversal or prevention of frailty in elderly humans, stimulation and

increase in muscle mass and muscle strength, attenuation of protein catabolic response after a major operation or trauma; reducing cachexia and protein loss due to chronic illness such as cancer or AIDS; improvement in muscle  
5 mobility, and maintenance of skin thickness.

As is well known to those skilled in the art, the known and potential uses of growth hormone are varied and multitudinous. Thus, the administration of the compounds of this invention for purposes of stimulating the release  
10 of endogenous growth hormone can have the same effects or uses as growth hormone itself. These varied uses of growth hormone may be summarized as follows: stimulating growth hormone release in elderly humans; maintenance of muscle strength and function in elderly humans, reversal  
15 or prevention of frailty in elderly humans, prevention of catabolic side effects of glucocorticoids, treatment of osteoporosis, stimulation and increase in muscle mass and muscle strength, stimulation of the immune system, acceleration of wound healing, acceleration of bone  
20 fracture repair, treatment of renal failure or insufficiency resulting in growth retardation, treatment of physiological short stature, including growth hormone deficient children, treatment of short stature associated with chronic illness, treatment of obesity and growth  
25 retardation associated with obesity, treating growth retardation associated with Prader-Willis syndrome and Turner's syndrome; accelerating the recovery and reducing hospitalization of burn patients; treatment of intrauterine growth retardation, skeletal dysplasia,  
30 hypercortisolism and Cushing's syndrome; induction of pulsatile growth hormone release; replacement of growth hormone in stressed patients; treatment of osteochondrodysplasias, Noonan's syndrome, schizophrenia, depression, Alzheimer's disease, delayed wound healing,  
35 and psychosocial deprivation; treatment of pulmonary dysfunction and ventilator dependency; attenuation of protein catabolic response after a major operation or

trauma; reducing cachexia and protein loss due to chronic illness such as cancer or AIDS; treatment of hyperinsulinemia including nesidioblastosis; adjuvant treatment for ovulation induction; to stimulate thymic development and prevent the age-related decline of thymic function; treatment of immunosuppressed patients; improvement in muscle mobility, maintenance of skin thickness, metabolic homeostasis, renal homeostasis in the frail elderly; stimulation of osteoblasts, bone remodeling, and cartilage growth; stimulation of the immune system in companion animals and treatment of disorders of aging in companion animals; growth promotant in livestock; and stimulation of wool growth in sheep.

In addition, the conditions, diseases, and maladies collectively referenced to as "Syndrome X" or Metabolic Syndrome as detailed in Johannsson *J. Clin. Endocrinol. Metab.*, 82, 727-34 (1997), may be treated employing the compounds of the invention.

The compounds of the present invention are agents that are growth hormone secretagogues and can be administered to various mammalian species, such as monkeys, dogs, cats, rats, humans, etc., in need of treatment. These agents can be administered systemically, such as orally or parenterally.

The compounds of the invention can be incorporated in a conventional systemic dosage form, such as a tablet, capsule, elixir or injectable formulation. The above dosage forms will also include the necessary physiologically acceptable carrier material, excipient, lubricant, buffer, antibacterial, bulking agent (such as mannitol), anti-oxidants (ascorbic acid or sodium bisulfite) or the like. Oral dosage forms are preferred, although parenteral intranasal or aerosol forms are quite satisfactory as well.

The dose administered must be carefully adjusted according to the age, weight, and condition of the patient, as well as the route of administration, dosage


form and regimen, and the desired result. In general, the dosage forms described above may be administered in amounts from about 0.0001 to about 100 mg/kg or body weight or in an amount within the range from about 1 to about 1000 mg per day, preferably, from about 5 to about 500 mg per day in single or divided doses of one to four times daily.

The following Examples represent preferred embodiments of the invention. All temperatures are in °C unless indicated otherwise.




#### GENERAL EXPERIMENTAL:

The term HPLCa refers to a Shimadzu high performance liquid chromatograph (HPLC) using a 4 minute gradient of from 0-100% solvent B [MeOH:H<sub>2</sub>O:0.2% H<sub>3</sub>PO<sub>4</sub>] with a 1 min. hold, an ultra violet (UV) detector set at 220nm, and using a column (4.6 x 50 mm) packed with YMC C18 5 micron resin. The term HPLCb refers to a Shimadzu HPLC using a 4 minute gradient of from 0-100% solvent B [MeOH:H<sub>2</sub>O:0.1% TFA], with a 1 min. hold, an ultra violet (UV) detector set at 220nm, and using a column (4.6 x 50 mm) packed with YMC C18 5 micron resin. A mixture of solvent A (10% MeOH/90% H<sub>2</sub>O/0.2% TFA) and solvent B (90% MeOH/10 %H<sub>2</sub>O/0.2% TFA) are used for preparative reverse phase HPLC in an automated Shimadzu system. The preparative columns are packed with YMC ODS C18 5 micron resin.

Example 1



A.



To a 0°C solution of the carboxylic acid

(5.0 g, 16.9 mmol, Sigma), dianiline

(2.3 g, 15 mmol, Aldrich), and 1-hydroxy-

zazabenzotriazole (HOAt) (2.5 g, 18 mmol) in DMF (48 ml)

under nitrogen was added 1-(3-dimethylaminopropyl)-3-

ethylcarbodiimide (EDAC or EDCI) (3.25 g, 17 mmol) then

diisopropylethyl amine (2.95 ml, 17 mmol). The reaction

mixture was allowed to slowly warm to room temperature

overnight. After 18 hrs, the reaction was diluted with

ethyl acetate. The organic layer was extracted with

dilute aqueous sodium bicarbonate, water, brine, dried

over sodium sulfate, and concentrated *in vacuo* to give

as a red colored oil (5.47 g),

containing starting dianiline, (referred to here as crude

compound). The above crude compound was used in the

subsequent reaction without further purification.

HPLCb rt = 3.19 min.

A portion of the above crude compound (1.25 g, ≤2.9

mmol) was dissolved in acetic acid (45 mL) and heated at

69°C for 4.5 h. After stirring at room temperature

overnight, the volatiles were removed *in vacuo* to give

crude title compound in the form of a red colored oil,

which may be used without further purification.


Alternatively, the red colored oil was dissolved in ethyl

acetate and the organic layers were extracted three times


with 1 N aqueous hydrogen chloride, water, brine, dried over sodium sulfate, and concentrated *in vacuo* to give pure title compound (0.92 g) as a dark colored oily foam. HPLCa rt = 3.99 min.

5

B.



Crude Part A compound ( $\leq 450$  mg,  $\leq 1.04$  mmol)) was treated with a solution of hydrogen chloride in dioxane (6 ml, 4N) and after 45 min. a brown solid had fallen from solution. The volatiles were evaporated *in vacuo* after 2 h. to give a dark colored solid. The residue was triturated three times with hot ethyl acetate to give



15 impure (crude) compound as a brown colored solid (338 mg). The crude compound was used without further purification in the next reaction. HPLCa rt = 2.93 min.

To a 0°C solution of Boc-methylalanine carboxylic acid (256 mg, 1.26 mmol, Aldrich), and the above crude compound (338 mg,  $\leq 0.97$  mmol), and HOAt (207 mg, 1.52 mmol) in DMF (5 ml) under nitrogen was added EDAC (245 mg, 1.28 mmol) then diisopropylethyl amine (0.39 ml, 2.25 mmol). The reaction mixture was allowed to slowly warm to room temperature overnight. After 18 h., the reaction was diluted with ethyl acetate. The organic layer was extracted with dilute aqueous sodium bicarbonate, 1N hydrogen chloride, water, brine, dried over sodium sulfate, and concentrated *in vacuo* to give a dark colored oil (564 mg). The crude residue was combined with material from another reaction (0.10 mmol) and was


30




purified by flash chromatography (SiO<sub>2</sub>, 40 g), eluting with 3.5% methanol/dichloromethane to give title compound as an orange colored oil (305 mg). HPLCb rt = 4.20 min; LC/MS (electrospray, + ions) m/z 498 (M+H).

5

C.



A solution of Part B compound (100 mg, 0.2 mmol) in methanol (2 ml) containing 10% palladium on carbon (15 mg) was purged with hydrogen and stirred under a hydrogen atmosphere (balloon) for 2 h. The reaction mixture was filtered through a Nylon frit, washing well with methanol, and the solvents evaporated to give amine compound




as an orange colored foam (82 mg, 88% crude recovery), ~80% pure by HPLCa rt = 3.0 min. The amine material was used without further purification in the next reaction. LC/MS (electrospray, + ions) m/z 468 (M+H).

To a solution of crude amine (82 mg, 0.18 mmol) and triethylamine (35  $\mu$ l, 0.25 mmol) in dichloromethane (1.8 ml) under nitrogen was added methanesulfonyl chloride (18  $\mu$ l, 0.23 mmol). After 2 h., the reaction was diluted with ethyl acetate and the organic layer was extracted with water, brine, dried over sodium sulfate, and concentrated in vacuo. The crude residue was purified by flash chromatography (SiO<sub>2</sub>, 10 g), eluting with 5.25% methanol/dichloromethane to give title compound as an orange colored solid (30.7 mg). HPLCa rt = 3.37 min.


30

D.




To neat Part C compound (31 mg, 56.8  $\mu$ mol) was  
5 added hydrogen chloride (0.8 ml, 4N in dioxane) and the  
mixture stirred for 2.5 h. The volatiles were removed in  
vacuo and co-evaporated twice with dichloromethane to give  
a tan colored solid. The crude residue was combined with  
material from another reaction (39.8  $\mu$ mol) and neutralized  
10 by dissolving in 9% methanol/dichloromethane containing  
ammonium hydroxide. The resulting suspension was purified  
by flash chromatography ( $\text{SiO}_2$ , 5 g), eluting with 9%  
methanol/dichloromethane containing 0.9% ammonium  
hydroxide to give a glassy solid (35 mg). The residue was  
15 dissolved in methanol (2.5 ml) and treated with  
trifluoroacetic acid (4.5  $\mu$ l). After 15 min, the  
volatiles were removed in vacuo to give the title compound  
(43 mg) as a tan colored solid. HPLCb rt = 2.43 min. MS  
(electrospray, + ions) m/z 446 (M+H).

20

Example 2


A.



25

*N*- Methyl morpholine (1.08 mL, 9.83 mmol) and isobutyl chloroformate (1.27 mL, 9.80 mmol) were added to a solution of *N*-Boc-O-benzyl serine (2.91 g, 9.84 mmol, Sigma) in THF (14.0 mL), cooled at -18°C. After stirring  
5 the mixture for 0.5 h. at -18°C, a -14°C solution of 3-aminopropionitrile (0.73 mL, 9.89 mmol) and *N*-methyl morpholine (1.08 mL, 9.83 mmol) in THF (6.0 mL) was added and the resulting mixture was allowed to warm up to -6°C over a period of 5.25 h. The mixture was filtered and the  
10 solution was evaporated near to dryness. The residue was taken up in EtOAc (70 mL) and washed with 5% NaHCO<sub>3</sub> (2x60 mL), water (60 mL) and brine (60 mL). The organic solution was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed(SiO<sub>2</sub> 230-400 mesh, 1/1 hexanes/EtOAc) to  
15 afford title compound (3.33 g, 97% yield): LC/MS (electrospray, + ions) m/z 348 (M+H).

B.




20

Diethylazodicarboxylate (1.51 mL, 9.59 mmol) and azidotrimethylsilane (1.27 mL, 9.57 mmol) were added to a solution of Part A compound (3.33 g, 9.60 mmol), triphenylphosphine (2.52 g, 9.61 mmol) and  
25 diisopropylethyl amine (0.42 mL, 2.41 mmol) in THF (30 mL), previously cooled at 0°C. The mixture was stirred at room temperature (rt) overnight and then cooled at 0°C. Another equivalent each of triphenylphosphine (2.52 g), diethylazodicarboxylate (1.51 mL) and  
30 azidotrimethylsilane (1.27 mL) was added and stirring was continued for an additional 28 h at rt. The reaction mixture was cooled at 0°C and mixed with an aqueous solution of ammonium cerium (IV) nitrate (2.63 g/100 mL). The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x150 mL)


and the combined organic phase was dried ( $\text{Na}_2\text{SO}_4$ ),  
 evaporated and chromatographed ( $\text{SiO}_2$  230-400 mesh, 1/1  
 hexanes/ $\text{EtOAc}$ ) to give the tetrazole (3.5 g, contaminated  
 with 1,2-dicarbethoxyhydrazine) and Part A compound (1.47  
 5 g contaminated with triphenylphosphine oxide). The  
 contaminated tetrazole was dissolved in  $\text{CHCl}_3$  and ether  
 was added to precipitate title compound (2.08 g) as a  
 colorless solid: MS (electrospray, + ions)  $m/z$  373 ( $M+H$ ).

10 C.



To a solution of Part B compound (354 mg, 0.95  
 mmol) in  $\text{CH}_2\text{Cl}_2$  (2.0 mL) was added a 4M  $\text{HCl}$ /dioxane  
 15 solution (3.0 mL) and the combined solution was allowed to  
 stand for 1.25h. The solvents were removed at reduce  
 pressure and the residue was taken up in *i*- $\text{PrOH}$  (1.5 mL).  
 Brine (25.0 mL) was added and the pH of the aqueous  
 solution was adjusted to 10 by addition of 1M  $\text{K}_2\text{CO}_3$ . The  
 20 aqueous solution was extracted with  $\text{CH}_2\text{Cl}_2$  (3x40 mL) and  
 the combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and  
 evaporated to provide title amine compound (258 mg) as a  
 colorless oil. This material was used without further  
 purification in the subsequent reaction:  $^1\text{H}$  NMR  $\delta$   
 25 ( $\text{CDCl}_3$ , ppm) 7.31 (m, 5H), 4.80 (m, 2H), 4.54 (m, 3H), 3.88  
 (m, 2H), 2.99 (m, 2H), 1.82 (broad s, 2H).


D.



30


CC(C(=O)O)C(=O)N (285 mg, 1.40 mmol, Sigma), EDAC (272 mg, 1.42 mmol), HOAt (190 mg, 1.40 mmol), 1,2-dichloroethane (1,2-DCE) (1.6 mL) and DMF (250  $\mu$ L) were mixed at 0°C and stirred for 15 min. The resulting cloudy solution was transferred, via syringe, to a 0°C solution of the crude Part C amine compound (258 mg, 0.95 mmol) in 1,2-DCE (1.4 mL) and the mixture was stirred for 20 h at rt. The reaction mixture was diluted with EtOAc (75 mL) and washed with saturated NaHCO<sub>3</sub> (2x40 mL), water (40 mL) and brine (40 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 1/1 hexanes/EtOAc) to provide title compound (338.5 mg) as a colorless oil: <sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>, ppm) 7.36 (m, 6H), 5.55 (m, 1H), 5.13 (broad s, 1H), 4.72 (t, *J*=6.6 Hz, 2H), 4.51 (2d, *J*=12 Hz, 2H), 4.02 (dd, *J*=9.2, 5.8 Hz, 1H), 3.86 (t, *J*=8.3 Hz, 1H), 2.98 (m, 2H), 1.44 (s, 3H), 1.41 (s, 3H), 1.32 (s, 9H).

E.




Part D compound (181 mg, 0.40 mmol) was treated with 15% TFA (4.0 mL) for 2.2 h and evaporated in vacuo and the residual TFA was coevaporated with CH<sub>2</sub>Cl<sub>2</sub> (5x6 mL). The oily crude residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>/MeOH (several drops) and ether was added to precipitate the title compound (157 mg) as a white solid: LC/MS (electrospray, + ions) m/z 358 (M+H).

## Example 3




A.



5

Diethylazodicarboxylate (1.51 mL, 9.59 mmol) and azidotrimethylsilane (1.27 mL, 9.57 mmol) were added to a solution of Example 2 Part A compound




10

(3.33 g, 9.60 mmol), triphenylphosphine (2.52 g, 9.61 mmol) and diisopropylethyl amine (0.42 mL, 2.41 mmol) in THF (30 mL), previously cooled at 0°C. The mixture was stirred at rt. overnight and then cooled at 0°C. Another equivalent each of triphenylphosphine (2.52 g), diethylazodicarboxylate (1.51 mL) and azidotrimethylsilane (1.27 mL) was added and stirring was continued for an additional 28 h at rt. The reaction mixture was cooled at 0°C and mixed with an aqueous solution of ammonium cerium (IV) nitrate (2.63 g/100 mL). The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x150 mL) and the combined organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 1/1 Hex/EtOAc) to give crude Example 2 Part B compound (3.5 g). The contaminated residue containing Example 2 Part B compound was dissolved in CHCl<sub>3</sub>, Example 2 Part B compound (2.08 g) precipitated by addition of ether, and the filtrate concentrated. The

25


resulting residue containing more Example 2 Part B compound was dissolved in THF (12 mL) and 1M NaOH and additional volumes of THF were added until disappearance of Example 2 Part B compound, monitoring by TLC (1/1  
5 hexanes/EtOAc). The final reaction mixture was diluted with brine (70 mL), and washed with ether (60 mL) and CHCl<sub>3</sub> (2x 50 mL). The aqueous layer pH was adjusted to 3, by addition of 1M H<sub>3</sub>PO<sub>4</sub>, and the solution was extracted with EtOAc (3x60 mL). The combined organic phase was  
10 washed with phosphate buffer (2x40 mL, pH 3) and brine (40 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to afford the the title compound (571 mg): LC/MS (electrospray, + ions) m/z 320 (M+H).

15 B.



Benzyl bromide (54  $\mu$ L, 0.45 mmol) was added to a stirred suspension of the Part A compound (115 mg, 0.36  
20 mmol) and cesium carbonate (258 mg, 0.79 mmol) in CH<sub>3</sub>CN (1.0 mL), and the mixture was stirred for 18 h. at rt. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and washed with water (8 mL). The aqueous phase was back extracted with CH<sub>2</sub>Cl<sub>2</sub> twice and the combined organic phase  
25 was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 3/1 hexanes/EtOAc) to give the title compound as a mixture of N-1/N-2 tetrazole isomers (122 mg): LC/MS (electrospray, + ions) m/z 410 (M+H).

C.




The isomeric mixture of Part B (122.2 mg, 0.30 mmol) was treated with 15% TFA (3.0 mL) for 2.0 h and evaporated. *i*-PrOH (2 mL) and brine (15.0 mL) were added to the residue and the pH of the aqueous solution was adjusted to 10 by addition of 1M K<sub>2</sub>CO<sub>3</sub>. The aqueous solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 25 mL) and the combined organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to provide title amine (94 mg). This amine material was used in the next reaction without further purification. LC/MS (electrospray, + ions) m/z 310 (M+H).

*N*-Boc-methyl alanine (79 mg, 0.39 mmol), EDAC (75 mg, 0.39 mmol), HOAT (53 mg, 0.40 mmol), 1,2-DCE (0.7 mL) and DMF (60  $\mu$ L) were mixed at 0°C and stirred for 15 min. The resulting cloudy solution was transferred, via syringe, to a 0°C solution of the crude amine (94 mg, 0.30 mmol) in 1,2-DCE (0.6 mL) and the mixture was stirred for 24 h at rt. The reaction mixture was diluted with EtOAc (30 mL) and washed with saturated NaHCO<sub>3</sub> (2x20 mL), water (20 mL) and brine (20 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 7/3 CHCl<sub>3</sub>/ether) to provide the title compound (44.6 mg). LC/MS (electrospray, + ions) m/z 495 (M+H).




D.




Part C compound (44.6 mg, 0.09 mmol) was treated  
 5 with ~2M HCl/CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (3.0 mL, prepared by  
 addition of AcCl to 8/1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) for 2.3 h. and  
 evaporated. The residue was further dried, under high  
 vacuum, to afford the title compound as a colorless solid  
 (41.1 mg): LC/MS (electrospray, + ions) m/z 395 (M+H).

10


Example 4

A.




15

Benzyl bromide (54  $\mu\text{L}$ , 0.45 mmol) was added to a  
 stirred suspension of the Example 3 Part A compound,




(115 mg, 0.36 mmol) and cesium carbonate (258 mg, 0.79 mmol) in CH<sub>3</sub>CN (1.0 mL), and the mixture was stirred for 18 h at rt. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and washed with water (8 mL). The aqueous phase was back extracted with CH<sub>2</sub>Cl<sub>2</sub> twice and the combined organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 3/1 hexanes/EtOAc) to give the title compound as a mixture of N-1/N-2 tetrazole isomers (122 mg): LC/MS (electrospray, + ions) m/z 410 (M+H).

B.



The isomeric mixture of Part A (122.2 mg, 0.30 mmol) was treated with 15% TFA (3.0 mL) for 2.0 h and evaporated. *i*-PrOH (2 mL) and brine (15.0 mL) were added to the residue and the pH of the aqueous solution was adjusted to 10 by addition of 1M K<sub>2</sub>CO<sub>3</sub>. The aqueous solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 25 mL) and the combined organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to provide




25

amine (94 mg). The material was used in the next reaction without further purification. LC/MS (electrospray, + ions)  $m/z$  310 (M+H).


5 *N*-Boc-methyl alanine (79 mg, 0.39 mmol), EDAC (75 mg, 0.39 mmol), HOAT (53 mg, 0.40 mmol), 1,2-DCE (0.7 mL) and DMF (60  $\mu$ L) were mixed at 0°C and stirred for 15 min. The resulting cloudy solution was transferred, via syringe, to a 0°C solution of the crude amine (94 mg, 0.30 mmol) in 1,2-DCE (0.6 mL) and the mixture was stirred for 10 24 h at rt. The reaction mixture was diluted with EtOAc (30 mL) and washed with saturated NaHCO<sub>3</sub> (2x20 mL), water (20 mL) and brine (20 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 7/3 CHCl<sub>3</sub>/ether) to provide the title compound (81.0 15 mg) as a white solid. LC/MS (electrospray, + ions)  $m/z$  495 (M+H).

C.




20

Part B compound (81.0 mg, 0.09 mmol) was treated with ~2M HCl/CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (3.0 mL, prepared by addition of AcCl to 8/1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) for 2.3 h. and evaporated. The residue was further dried, under high 25 vacuum, to afford the title compound (72.4 mg) as a pale yellow solid: LC/MS (electrospray, + ions)  $m/z$  395 (M+H).


Example 5

A.




5

1M NaOH (0.67 mL) and *i*-PrOH (0.50 mL) were added to a solution of Example 2 Part D compound




- 10 (308 mg, 0.67 mmol) in 6/1 THF/Dioxane (3.5 mL), and the mixture was stirred at rt. for 13 h. The reaction mixture was diluted with brine (30 mL) and the pH of the aqueous solution was adjusted to 3 by addition of 1M H<sub>3</sub>PO<sub>4</sub>. The aqueous solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x50 mL) and
- 15 the combined organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to yield the title compound (234 mg): LC/MS (electrospray, + ions) m/z 405 (M+H).

B.



Diisopropylethyl amine (10  $\mu$ L, 57  $\mu$ mol) and  
 5 diethylazodicarboxylate (53  $\mu$ L, 0.34 mmol) were added to a  
 solution of (+)-methyl-L-mandelate (56 mg, 0.34 mmol,  
 Aldrich), Part A compound (118.5 mg, 0.29 mmol) and  
 triphenylphosphine (87 mg, 0.33 mmol) in  $\text{CH}_2\text{Cl}_2$  (0.9 mL),  
 previously cooled at  $0^\circ\text{C}$ . After stirring the mixture at  
 10 rt for 34 h, additional amounts of the reagents,  
 triphenylphosphine (99 mg), of (+)-methyl-L-mandelate (56  
 mg) and diethylazodicarboxylate (53  $\mu$ L), were added. The  
 mixture was stirred at rt for an additional 24 h and  
 concentrated *in vacuo*. The resulting residue was  
 15 chromatographed ( $\text{SiO}_2$  230-400 mesh, 4/1  $\text{CH}_2\text{Cl}_2$ /Ether) to  
 afford a mixture containing the title compound (57.5 mg)  
 contaminated with 1,2-dicarbethoxyhydrazide. The mixture  
 was purified by chromatography ( $\text{SiO}_2$  230-400 mesh, 95/5  
 $\text{CHCl}_3$ /MeOH) to provide the title compound (49.2 mg, 1/1  
 20 mixture of diastereomers) as a colorless oil: LC/MS  
 (electrospray, + ions)  $m/z$  553 ( $M+H$ ).


C.



25


Part B compound (49.2 mg, 89  $\mu$ mol) was treated with  
 ~2M HCl/CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (3.0 mL, prepared by addition  
 of AcCl to 8/1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) for 2.3 h and evaporated. The  
 residue was further dried, under high vacuum, to afford  
 5 the title compound (42.7 mg, 1/1 mixture of diastereomers)  
 as a white solid: LC/MS (electrospray, + ions) m/z 453  
 (M+H).

Example 6




10

A.




15 Diisopropylethyl amine (10  $\mu$ L, 57  $\mu$ mol) and  
 diethylazodicarboxylate (53  $\mu$ L, 0.34 mmol) were added to a  
 solution of (+)-methyl-L-mandelate (56 mg, 0.34 mmol,  
 Aldrich), compound Example 5 Part A



20 (118.5 mg, 0.29 mmol) and triphenylphosphine (87 mg, 0.33  
 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.9 mL), previously cooled at 0°C. After  
 stirring the mixture at rt for 34 h, additional amounts of

the reagents, triphenylphosphine (99 mg), of (+)-methyl-L-mandelate (56 mg) and diethylazodicarboxylate (53  $\mu$ L), were added. The mixture was stirred at rt for an additional 24 h and concentrated *in vacuo*. The resulting residue was chromatographed (SiO<sub>2</sub> 230-400 mesh, 4/1 CH<sub>2</sub>Cl<sub>2</sub>/Eter) to afford the title compound (70.6 mg, 1/1 mixture of diastereomers) as a colorless oil: LC/MS (electrospray, + ions) m/z 553 (M+H).


10 B.




Part A compound (70.6 mg, 0.13 mmol) was treated with ~2M HCl/CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (3.0 mL, prepared by addition of AcCl to 8/1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) for 2.5 h and evaporated. The residue was further dried, under high vacuum, to afford the title compound (61.9 mg, 1/1 mixture of diastereomers) as a white solid: LC/MS (electrospray, + ions) m/z 453 (M+H).

20

#### Example 7




A.




25

To a precooled 160 ml of methanol at 0°C was added acetyl chloride (10g, 128 mmol) slowly. After stirring at 0°C for 15 min, O-benzyl-D-serine




- 5 (5g, 25.6 mmol) was added and the solution was heated to reflux for 2 days. The solvent was removed and the residue was dried under high vacuum to give title compound as a white solid (6.29g, 100%). HPLCa rt = 1.90 min.

10 B.



- To a solution of Part A compound (3.65g, 14.9 mmol) in 100 ml of dry methylene chloride was added di-t-butylcarboxymate (3.89g, 17.8 mmol) followed by the addition of triethyl amine (2.25g, 22.3 mmol). The reaction was stirred overnight and quenched with 50 ml of water and extracted with methylene chloride (3x100ml). The combined organic layers were washed with brine, dried over Na2SO4, and concentrated to give crude title product as an oil (5.01g). HPLCb rt=3.24 min.; LC/MS (electrospray, + ions) m/z 310 (M+H).

C.



25


To a solution of crude Part B compound (0.98g, 3.17 mmol) in 10 ml of dry methyl alcohol was added hydrazine (0.447g, 13.9 mmol). The reaction was stirred at rt for 3



h and heated to reflux for 5 h. The solvent was removed and the residue was dried under high vacuum to give title compound as an oil (0.96g, 98%). HPLCb rt = 3.21 min.; LC/MS (electrospray, + ions) m/z 310 (M+H).

5


D.



To a solution of crude Part C compound (0.53 g, 1.72 mmol) in 10 ml of dry ethyl alcohol was added benzyl isothiocyanate (307 mg, 2.06 mmol). The reaction was heated to reflux for 4 h. The reaction was allowed to cool to room temperature and the solvent was removed in vacuo. The residue was then treated with 1N NaOH and the resulting mixture was heated to 109°C for 4 h. After cooling to rt, the resulting white precipitate was isolated by filtration. The solid was rinsed with water and dried under high vacuum to give title compound as a white solid (0.36g, 62%). HPLCb rt = 3.29 min.; LC/MS (electrospray, + ions) m/z 341 (M+H).

20

E.




To a solution of Part D compound (153 mg, 0.45 mmol) in 4 ml of DMF was added Boc-methylalanine carboxylic acid (96 mg, 0.47 mmol, Sigma), HOAt (64 mg, 0.47 mmol), EDAC (91 mg, 0.47 mmol), and diisopropylethyl amine (61 mg, 0.47 mmol) at rt under nitrogen. The reaction mixture was stirred at rt for 20 h, and quenched with water (10 ml). The mixture was extracted with ethyl

30

acetate (3x20 ml), the organic layer dried over sodium sulfate, and concentrated in vacuo to give a crude yellow oil. The crude residue was purified by flash chromatography, eluting with hexane/ethyl acetate (2:1) to give title compound as an oil (153 mg, 65%). HPLCb rt = 4.13 min.; LC/MS (electrospray, + ions) m/z 526 (M+H).


F.



10


Part E compound (133 mg, 0.25 mmol) was treated with 2 ml of TFA/CH<sub>2</sub>Cl<sub>2</sub> (1:3) at room temperature for 1 h, the solvent removed and the residue was purified by preparative HPLC (Shimadzu, 30-100% B, 30 min. gradient, 5 min. hold, 220nM) to give the title compound (80 mg, 59%). HPLCb rt = 3.25 min.; LC/MS (electrospray, + ions) m/z 425 (M+H).

#### Example 8




20


Example 7 Part E, compound (133 mg, 0.25 mmol) was treated with 2 ml of TFA/CH<sub>2</sub>Cl<sub>2</sub> (1:3) at room temperature for 1 h, the solvent removed and the residue was purified by preparative HPLC (Shimadzu, 30-100% B, 30 min. gradient, 5 min. hold, 220nM) to give the title compound (42 mg, 28%). HPLCb rt = 3.80 min.; LC/MS (electrospray, + ions) m/z 481 (M+H).

Example 9

A solution of Example 7 Part E, compound (78 mg, 0.15 mmol) in 7.5 ml of THF was added to hydrogen peroxide (0.33 ml, 30% w/w). After stirring at rt for 2 days, the solvent was removed. The residue was then treated with 3 ml of TFA/CH<sub>2</sub>Cl<sub>2</sub> (1:3) at room temperature for 1 h, the solvent removed, and the residue was purified by preparative HPLC (Shimadzu, 20-100% B, 30 min. gradient, 5 min. hold, 220nm) to give the title compound (59 mg, 79%). HPLCb rt = 2.99 min.; LC/MS (electrospray, + ions) m/z 394 (M+H).

Example 10

A.





20

To a suspension of imidazole (19.08 g, 0.28 mol) in 300 ml of dry benzene was added dimethylchlorosulphonamide (26 ml, 0.24 mol) and triethylamine (36 ml, 0.26 mol). The reaction was stirred at rt under nitrogen for 24 h, the mixture filtered, and the filtrate was concentrated. The resulting crude product was distilled under reduced pressure (0.03 mmHg) to yield title compound (36.4g, 86%). HPLCb rt = 0.64 min.; LC/MS (electrospray, + ions) m/z 176 (M+H).


30

B.



To a suspension of Part A compound (1.88 g, 10.75  
 5 mmol) in 100 ml of dry THF at  $-78^{\circ}\text{C}$  was added n-BuLi  
 (4.60 ml, 2.5 M in hexane, 11.51 mmol) dropwise under  
 nitrogen. The reaction was stirred at  $-78^{\circ}\text{C}$  for 1 h and  
 4-phenyl-butylaldehyde  in 20 ml of dry ether  
 was added in one portion. The mixture was stirred at -  
 10  $78^{\circ}\text{C}$  for 5 min. and then allowed to go to rt. After 3 hr  
 at rt, the reaction was quenched with sat.  $\text{NH}_4\text{Cl}$  solution.  
 The mixture was extracted with ethyl acetate (3x20 ml),  
 the organic layer dried over sodium sulfate, and  
 concentrated *in vacuo* to give a crude yellow oil. The  
 15 crude residue was purified by flash chromatography,  
 eluting with hexane/ethyl acetate (1:3) to give title  
 compound as an oil (3.24 g, 93%). HPLCb rt = 3.44 min.;  
 LC/MS (electrospray, + ions) m/z 324 (M+H).


20 C.



To a mixture of Part B compound (3.2 g, 9.91 mmol)  
 and diphenyl phosphorazide (3.27 g, 11.89 mmol) in 17 ml  
 25 of dry toluene at  $0^{\circ}\text{C}$  under nitrogen was added 1,8-  
 diazabicyclo[5.4.0]undec-7-ene (DBU) (1.81 g, 11.89 mmol).  
 The reaction was stirred for 2 h at  $0^{\circ}\text{C}$  and then 20 h at  
 room temperature. The resulting two phase mixture was  
 quenched with water and extracted with ethyl acetate (3x20  
 30 ml). The organic layer was washed with brine, dried over  
 sodium sulfate, concentrated *in vacuo* and purified by  
 flash chromatography using 3:1 hexane/ethyl acetate to

afford title compound as an oil (1.64 g, 48%). HPLCb rt = 4.15 min.; LC/MS (electrospray, + ions) m/z 349 (M+H).


D.



5

A mixture of Part C compound (232 mg, 0.67 mmol) and 5% Pd/C (50 mg) in 3 ml of EtOH was hydrogenated under 1 atmosphere of hydrogen. After 9 h, the solid was then  
10 filtered off and the filtrate was concentrated in vacuo to afford Part D compound as an oil (200 mg, 93%). HPLCb rt = 3.10 min.; LC/MS (electrospray, + ions) m/z 323 (M+H).


E.



15

To a solution of Part D compound (172 mg, 0.53 mmol) and Boc-methylalanine carboxylic acid (114 mg, 0.56 mmol, Sigma) in 3 ml of acetonitrile was added  
20 benzotriazol-1-yloxy-bis(dimethylamino)phosphonium hexafluorophosphate (BOP) (248 mg, 0.56 mmol). After stirring for 20 min at rt, triethylamine (56.6 mg, 0.56 mmol) was added and the reaction was stirred for 5 h at room temperature. The solvent was removed via evaporation  
25 and the residue was purified by flash chromatography, eluting with hexane/ethyl acetate (1:1) to give Part E compound as an oil (238 mg, 88%). HPLCb rt = 4.13 min.; LC/MS (electrospray, + ions) m/z 508 (M+H).








F.




Part E compound (167 mg, 0.33 mmol) was treated  
5 with 2 ml of TFA/CH<sub>2</sub>Cl<sub>2</sub> (1:3) at room temperature for 1 h,  
the solvent was removed and the residue was purified by  
preparative HPLC (Shimadzu, 30-100% B, 30 min. gradient, 5  
min. hold, 220nm) to give the title compound (137 mg).  
HPLCb rt = 3.33 min.; LC/MS (electrospray, + ions) m/z 408  
10 (M+H).


The following compounds were prepared employing the  
procedures described above and the working Examples.


Examples 11 to 45

Example No.	Structure	M+H positive ions
11	 Chiral	387
12	 Chiral	319
13	 Chiral	421
14	 Chiral	421
15	 Chiral	440
16	 Chiral	440
17	 Chiral	488











44



Chiral

494


45




Chiral

570

## Example 46



A.




5

N-Methyl morpholine (2.5 mL) and isobutyl  
 chloroformate (2.88 mL, 22.8 mmol) were added to a  
 solution of N-Boc-O-benzyl-D-serine (6.73 g, 22.8 mmol,  
 10 Sigma) in THF (35 mL), cooled at  $-18^{\circ}\text{C}$ . After stirring  
 the mixture for 0.5 h. at  $-18^{\circ}\text{C}$ , a THF (15 mL) and DMF (25  
 mL) solution of methyl 4-aminobutyrate hydrochloride  
 $\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CO}_2\text{Me}$  (3.53 g, 23 mmol) and N-methyl morpholine  
 (5 mL) was added, keeping the reaction mixture below  
 15  $-10^{\circ}\text{C}$ . The resulting mixture was stirred for 5 h. at  
 between  $-10^{\circ}$  to  $-15^{\circ}\text{C}$ . The mixture was filtered, washing  
 with ethyl acetate, and the solution was evaporated under  
 reduced pressure. The residue was taken up in EtOAc (160  
 mL) and washed with 5%  $\text{NaHCO}_3$  (2x40 mL), water (50 mL) and  
 20 brine (50 mL). The organic solution was dried ( $\text{MgSO}_4$ ),  
 evaporated and chromatographed ( $\text{SiO}_2$  230-400 mesh, 4%  
 methanol/ dichloromethane) to afford title compound (8.5  
 g) as a colorless syrup: LC/MS (electrospray, + ions) m/z  
 395 (M+H).


25

B.



Diethylazodicarboxylate (13.4 mL, 21.6 mmol) and  
5 azidotrimethylsilane (2.87 mL, 21.6 mmol) were added to a  
0°C solution of Part A compound (8.5 g, 21.6 mmol),  
triphenylphosphine (5.7 g, 21.6 mmol) and diisopropylethyl  
amine (1 mL, 5.4 mmol) in THF (70 mL). The mixture was  
stirred at rt. for 24 h., and then cooled at 0°C. Another  
10 equivalent each of triphenylphosphine (5.7 g),  
diethylazodicarboxylate (13.4 mL) and azidotrimethylsilane  
(2.87 mL) was added and stirring was continued for an  
additional 48 h at rt. The reaction mixture was then  
concentrated in vacuo to one third the original volume,  
15 diluted with ethyl acetate (200 mL), then cooled to 0°C.  
An aqueous solution of ammonium cerium (IV) nitrate (65  
g/250 mL) was added and the mixture stirred for 15 min.  
The organic layer was separated, the aqueous mixture was  
extracted with ethyl acetate (150 mL), and the combined  
20 organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>). After concentration in  
vacuo, the residue was chromatographed (SiO<sub>2</sub> 230-400 mesh,  
40% ethyl acetate/hexanes) to give solid title compound  
(5.5 g, contaminated with a less polar impurity) and  
recovered Part A compound (5.5 g, contaminated with  
25 triphenylphosphine oxide). The contaminated title  
compound could be used as is in the subsequent reaction.  
Alternatively, a portion of the impure title compound (4.2  
g) was chromatographed (SiO<sub>2</sub> 230-400 mesh, 25% ethyl  
ether/dichloromethane) to give solid title compound (3.5  
30 g) as a colorless solid: LC/MS (electrospray, + ions) m/z  
420 (M+H).


C.



To a solution of Part B compound (3.5 g, 8.48 mmol) in dichloromethane (20 mL) was added a 4M HCl/dioxane solution (5 mL). After 3 h., more 4M HCl/dioxane solution (4 mL) was added and after a total of 5 h., the volatiles were removed in vacuo to give a syrup. The residue was dissolved in dichloromethane (200 mL), washed with 1N NaOH (75 mL), brine (2x80 mL), dried (sodium sulfate), and concentrated to give title compound (2.46 g) as a syrup. This material was used without further purification in the subsequent reaction: LC/MS (electrospray, + ions) m/z 320 (M+H).

15


D.



A DMF (30 mL) solution of *N*-Boc-methyl alanine (1.88 g, 9.2 mmol, Sigma), Part C compound (2.46 g, 7.7 mmol), HOAT (1.17 g, 8.5 mmol), and EDAC (1.62 g, 8.5 mmol) was stirred at room temperature for 20 h. The reaction mixture was concentrated in vacuo and the residue dissolved in dichloromethane (200 mL). The organic layer was washed with water (3X100 mL), dried over magnesium sulfate, and concentrated in vacuo. The residue was chromatographed (SiO<sub>2</sub> 230-400 mesh) eluting with 40% ether/dichloromethane to give title compound (3.14 g) as a colorless oil: LC/MS (electrospray, + ions) m/z 505 (M+H).


30

E.




To a solution of Part D compound (1.4 g, 2.78 mmol)  
 5 in THF (8 mL) was added an aqueous solution of lithium  
 hydroxide (143 mg, 6 mmol, in 2 mL water) and the mixture  
 was allowed to stir for 20 h. at room temperature. The  
 reaction mixture was brought to pH 1-2 with 1N aq.HCl (5  
 mL) and the aqueous layer was extracted with ethyl acetate  
 10 (200 mL). The organics were washed with water (60 mL),  
 brine (100 mL), dried (magnesium sulfate), and  
 concentrated in vacuo to give title compound (1.32 g) as a  
 colorless solid. This material was used without further  
 purification in the subsequent reaction: LC/MS  
 15 (electrospray, + ions) m/z 491 (M+H).

F.




20 To tryptamine



(24 mg, 0.15 mmol) was added a dichloromethane (1 mL) and  
 DMF (0.5 mL) solution of Part E compound (50 mg, 0.1  
 mmol), EDAC (28.8 mg, 0.15 mmol), and dimethylamino  
 25 pyridine (18.3 mg, 0.15 mmol, DMAP). The reaction mixture  
 was shaken for 20 h., then diluted with methanol (5 mL).

The resulting solution was passed through a SCX resin column (1 g), eluting with methanol (15 mL), and the eluent was concentrated in vacuo to give the crude intermediate amide:

5




The resulting residue was dissolved in a dichloromethane/methanol HCl solution (1.5 mL, prepared from dissolving acetyl chloride (10.2 mL) in 3/2 dichloromethane/methanol  
10 (40 mL), 3.5 M) and stirred at rt. for 4 h. before removing the volatiles under reduced pressure. The resulting residue was purified by Prep HPLC to give the title compound as a dark colored foam (44 mg): LC/MS (electrospray, + ions) m/z 533 (M+H).














15 In a manner analogous to that of preparing the Example 46 compound, compounds of Examples 47 to 91 listed in the table below were prepared from Example 46 Part E compound (0.1 mmol) and the respective amine (0.15 to 0.2 mmol). Examples 47 to 50 were treated with trifluoroacetic  
20 acid (0.4 mL) and thioanisole (0.05 mL) in dichloromethane (1 mL) instead of HCl to form the final products. All but Example 70 were purified by preparative HPLC, eluting with a gradient system of methanol and water with 0.2% trifluoroacetic acid. All compounds were isolated as  
25 trifluoroacetic acid salts.


















In the tables of compounds which follow, the X<sub>1</sub> designation refers to the point of attachment of the particular R moiety shown to the rest of the molecule.




Examples 47 to 91

Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
47		508
48		472
49		480
50		494
51		446
52		460
53		504
54		490
55		476
56		462
57		448

58		481
59		495
60		481
61		506
62		466
63		510
64		545
65		558
66		510
67		497
68		486
69		573
70		447


71		462
72		443
73		476
74		434
75		476
76		478
77		494
78		448
79		462
80		490
81		475
82		502
83		515
84		503
85		464
86		478
87		496

88




510

89




526

90




508

91




462


Example 92

5

A.



Methyl triflate (1.30 mL, 11.5 mmol) was added to a suspension of resin bound (tert-alkoxycarbonyl)imidazole




- 10 (8.68 g, loading 0.72 mmol/g) [Note: this procedure was adapted from Hernandez and Hodges. *J. Org. Chem.* 1997, 62, 3153] in dry 1,2-DCE (60.0 mL), cooled at 5°C. The mixture was stirred for 20 min at this temperature and for 15 min while being warmed to rt.
- 15 After addition of DMAP (2.96 g, 24.2 mmol), stirring was continued for an additional 10 min.

20 *N,O*-Bis(trimethylsilyl)acetamide (4.48 mL, 18.1 mmol) was added to a suspension of *O*-benzyl-D-serine (3.54 g, 18.1 mmol) in DMF (21 mL) and the mixture was stirred for 30 min at rt. The resulting solution was transferred

via syringe to the stirred resin suspension. The mixture was shaken for 8 h at rt and filtered. The resin bound *O*-benzyl-D-serine was washed with DMF (3 times), 5% AcOH/DMF, MeOH (3 times, 100 mL total), THF (3 times), and CH<sub>2</sub>Cl<sub>2</sub> (3 times) and dried: 9.58 g; IR 1717 (broad) cm<sup>-1</sup>; Anal. Found C, 80.78; H, 8.00; N, 0.91. Loading on N content: 0.65 mmol/g.

A portion of the resin (27 mg) was treated with 10% TFA/ CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) for 4.5 h and filtered. The resin was rinsed with CH<sub>2</sub>Cl<sub>2</sub> (3 times) and MeOH (2 times) and the filtrates were evaporated and dried (vacuum, overnight) to give back *O*-benzyl-D-serine, as its TFA salt (4.4 mg): LC-MS 100% Area; LC/MS (electrospray, + ions) m/z 196 (M+H). Loading, on cleaved amount, of the resin bound *O*-benzyl-D-serine: 0.53 mmol/g.

B.




To the Part A resin bound *O*-benzyl-D-serine (1.75 g, 1.14 mmol) was added a solution of β-alanine methyl ester hydrochloride (315 mg, 2.27 mmol), diisopropylethylamine (0.39 mL, 2.24 mmol) and HOAT (308 mg, 2.26 mmol) in DMF (3.0 mL) and then, a solution of EDAC (433 mg, 2.26 mmol) in DMF (5.5 mL). The mixture was rocked for 24 h at rt and filtered. The polymer was washed 3 times each with DMF, THF and CH<sub>2</sub>Cl<sub>2</sub> to give resin bound *O*-benzyl-D-serine-2-methoxy-carbonylethyl amide.

To the resin bound amide was added a solution of triphenylphosphine (955 mg, 3.64 mmol) in 1,2-DCE (3.0 mL) and a solution, previously cooled at 0°C, of trimethylsilylazide (0.48 mL, 3.62 mmol) and diethyl azodicarboxylate (DEAD) (0.57 mL, 3.62 mmol) in 1,2-DCE (3.0 mL). The mixture was rocked for 24 h at rt and filtered. The resin was washed with DMF (3 times) and

CH<sub>2</sub>Cl<sub>2</sub> (6 times), and resubmitted to the above Mitsunobu reaction conditions twice. The polymer was finally washed with DMF (3 times), THF (3 times) and CH<sub>2</sub>Cl<sub>2</sub> (3 times) to provide the resin bound 1-(2-methoxycarbonylethyl)-  
 5 tetrazole.

C.




10 To the Part B resin bound methyl ester resin (307 mg, < 0.20 mmol) in *n*-butyl amine (2.0 mL) was added a solution of tetra-*n*-butyl ammonium cyanide (40 mg, 0.15 mmol) in MeOH (0.3 mL) and the mixture was heated at 50°C for 3.6 h. After filtration, the polymer was washed 3  
 15 times each with DMF, MeOH, THF and CH<sub>2</sub>Cl<sub>2</sub> to give the resin bound amide.

The resin was treated with ~3M HCl/ CH<sub>2</sub>Cl<sub>2</sub>, MeOH, methyl acetate (4.0 mL, prepared by addition of acetyl chloride (AcCl) to a 3/2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH solution) for 4.5 h  
 20 and filtered. The resin was rinsed with CH<sub>2</sub>Cl<sub>2</sub> (3 times) and MeOH (twice), and the filtrates were evaporated to afford the amine HCl salt intermediate. This material was taken up in isopropanol (1.5 mL) and saturated NaCl (20 mL). The pH of the aqueous phase was adjusted to 10 by  
 25 addition of 1M K<sub>2</sub>CO<sub>3</sub> and the solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x30 mL). The combined organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give the free amine contaminated with triphenylphosphine oxide (30.7 mg).

N-Boc isobutyric acid (36 mg, 0.18 mmol), EDAC (34 mg, 0.18 mmol), HOAT (25 mg, 0.18 mmol), 1,2-DCE (0.3 mL) and DMF (30 uL) were mixed at 0°C for 15 min. The  
 30 resulting solution was added to a 0°C solution of the


crude amine (30.7 mg, < 0.09 mmol) in 1,2-DCE (0.2 mL). The mixture was stirred for 22 h at rt and diluted with EtOAc (25 mL). The solution was washed with saturated NaHCO<sub>3</sub> (2x20 mL), water (20 mL) and saturated NaCl (20 mL), dried and evaporated to give crude Part C compound (44.4 mg). Preparative HPLC (solvent B: start 30%, final 90%; gradient time: 15 min; flow rate 20 mL/min; wavelength 220 nm; column YMC S5 ODS 20x100 mm) gave the Part C compound (19.0 mg): LC/MS (electrospray, + ions) m/z 532 (M+H).

D.




The Part C compound (19.0 mg, 0.036 mmol) was treated with ~3M HCl in CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (2.0 mL, prepared by addition of AcCl to a 3/2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH solution) for 2.2 h and concentrated. The residue was further dried (vacuum, 2 h) to afford the title compound (16.6 mg) as a colorless solid: LC-MS 96% Area; LC/MS (electrospray, + ions) m/z 432 (M+H).


#### Example 93



A.




- 5 Methyl triflate (0.80 mL, 7.11 mmol) was added to a suspension of resin bound (tert-alkoxycarbonyl)imidazole (5.06 g, Loading 0.70 mmol/g)



- 10 in dry 1,2-DCE (30 mL), cooled at 10°C. The mixture was stirred for 15 min at this temperature and for 10 min while being warmed to rt. After addition of Et<sub>3</sub>N (2.40 mL, 17.2 mmol), stirring was continued for an additional 10 min. A suspension of methyl 2-aminoisobutyrate (1.14 g, 7.42 mmol) and Et<sub>3</sub>N (0.98 mL, 7.04 mmol) in DMF (13 mL)
- 15 was filtered and transferred via syringe to the stirred resin suspension. The mixture was shaken for 5.5 h at rt and filtered. The resin bound Part A compound was washed with THF (3 times), 1/1 THF/MeOH, THF (3 times), and CH<sub>2</sub>Cl<sub>2</sub> (3 times) and dried: IR 1728 cm<sup>-1</sup>; Anal. Found C, 82.02; H, 7.96; N, 0.89. Loading on N content: 0.64
- 20 mmol/g.

B.



25

- A 3/2 dioxane/0.25 M KOH solution (40 mL) was added to the Part A compound (2.63 g, Loading 0.64 mmol/g) and the suspension was heated at 75°C for 4.5
- 30 h. After filtration, the polymer was washed with DMF (3 times), 5% AcOH/DMF (3 times, 50 mL total), MeOH (3




times), THF (3 times) and  $\text{CH}_2\text{Cl}_2$  (3 times) and dried to provide the Part B compound (2.50 g).

A portion of the resin (53 mg) was treated with 10% TFA/  $\text{CH}_2\text{Cl}_2$  (1.5 mL) for 5 h and filtered. The resin was rinsed with  $\text{CH}_2\text{Cl}_2$  (3 times) and MeOH (2 times) and the filtrates were evaporated and dried (vacuum, overnight) to give pure 2-aminoisobutyric acid, as its TFA salt (6.3 mg):  $^1\text{H}$  NMR  $\delta$  ( $\text{CD}_3\text{OD}$ , ppm) 1.55 (s, 6H).

Loading, on cleaved amount, of the resin bound 2-aminoisobutyric acid Part B compound: 0.53 mmol/g.


C.



15

To a suspension of Example 92, Part B compound in phenethylamine (1.4.0 mL) was added a solution of tetra-*n*-butyl ammonium cyanide (22 mg, 0.08 mmol) in MeOH (0.3 mL) and the mixture was heated at 60-65°C for 3 h. After filtration, the polymer was washed 3 times each with DMF, 1/1 THF/MeOH, THF and  $\text{CH}_2\text{Cl}_2$  to give the resin bound amide.


The resin was treated with ~3M HCl in  $\text{CH}_2\text{Cl}_2$ , MeOH, MeOAc (3.0 mL, prepared by addition of AcCl to a 3/2  $\text{CH}_2\text{Cl}_2$ /MeOH solution) for 5 h and filtered. The resin was rinsed with  $\text{CH}_2\text{Cl}_2$  (3 times) and MeOH (twice), and the filtrates were evaporated to afford the intermediate



(69.3 mg).


The above intermediate (69.3 mg, 0.18 mmol), Part A compound (400 mg, loading 0.53 mmol/g) and a solution of EDAC (115 mg, 0.60 mmol), HOAT (82 mg, 0.60 mmol) and diisopropylethylamine (0.10 mL, 0.60 mmol) in DMF (3.0 mL) were mixed and the resulting suspension was rocked at rt for 24 h. After filtration, the resin was washed with DMF (3 times) and dioxane (3 times). The resin was swollen with 3/2 dioxane/H<sub>2</sub>O, rocked overnight, filtered and washed with THF (3 times) and CH<sub>2</sub>Cl<sub>2</sub> (3 times).

The resin was treated with ~3M HCl in CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeOAc (4.0 mL, prepared by addition of AcCl to a 3/2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH solution) for 6 h and filtered. The resin was rinsed with CH<sub>2</sub>Cl<sub>2</sub> (2 times) and MeOH and the filtrates were evaporated to give the crude title compound. This crude mixture was dissolved in acetonitrile/MeOH and passed through an anion exchange cartridge (3.0 g, CHQAX) by eluting with acetonitrile (18 mL) to provide the amine (22.2 mg). Further elution with 1/1 acetonitrile/MeOH (20 mL) gave an additional amount of amine (29.8 mg). The combined amine was further purified by preparative HPLC (solvent B: start 30%, final 80%; gradient time: 20 min; flow rate 20 mL/min; wavelength 220 nm; column YMC S5 ODS 20x100 mm) to afford the title compound (20.7 mg): HPLC 100% Area; LC/MS (electrospray, + ions) m/z 480 (M+H).

Examples 94 and 95


5

A.




- Sodium borohydride (110 mg, 2.92 mmol) was added to a solution of cinnamoyl acetonitrile (1.0 g, 5.84 mmol, Maybridge) in 4/1 MeOH/H<sub>2</sub>O (30 mL), cooled at 0°C, and the mixture was stirred for 1.5 h. After warming to rt, small portions of sodium borohydride were added at 45 min intervals until reaction completion (additional 90 min). After evaporating *in vacuo* most of the MeOH, potassium phosphate buffer (pH 3, 45 mL) was added and the mixture was stirred for 15 min. 5% NaHCO<sub>3</sub> (45 mL) was added and the aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic phases were combined, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was chromatographed (SiO<sub>2</sub> 230-400 mesh, 1/1 Hex/EtOAc) to give the Part A compound (842.5 mg) as a yellow liquid: LC/MS 98% Area; LC/MS (electrospray, + ions) m/z 156 (M-OH).


B.




Diisopropylethylamine (31  $\mu$ L, 0.18 mol) and  
 5 diethylazodicarboxylate (0.18 mL, 1.14 mmol) were added to  
 a solution of the Part A alcohol (198 mg, 1.14 mmol),  
 Example 5 Part A compound



(360 mg, 0.89mmol) and  $\text{Ph}_3\text{P}$  (303 mg, 1.15 mmol) in  $\text{CH}_2\text{Cl}_2$   
 10 (2.6 mL), previously cooled at  $0^\circ\text{C}$ . The mixture was  
 stirred at rt for 24 h and evaporated. The residue was  
 chromatographed ( $\text{SiO}_2$  230-400 mesh, 1/1 Hex/EtOAc) to  
 afford the Part B N-1 substituted tetrazoles (98.3 mg, 20%  
 yield, 2 diastereomers): HPLC 98% diastereomer combined  
 15 Area; LC/MS (electrospray, + ions) m/z 560 (M+H).



and



20

A solution of the Part B compound (72 mg, 0.13  
 mmol, 2 diastereomers) and thioanisole (45  $\mu$ L, 0.38 mmol)  
 in  $\text{CH}_2\text{Cl}_2$  (1.5 mL) was treated with 4M HCl/dioxane  
 solution (1.5 mL) for 1.2 h and concentrated. Preparative






HPLC (solvent B: start 40%, final 90%; gradient time: 30 min; flow rate 20 mL/min; wavelength 220 nm; column YMC S5 ODS 20x100 mm) afforded Example 94 title compound (23.8 mg, 32% yield, single diastereomer) as a colorless solid:

5 LC/MS 100% Area; LC/MS (electrospray, + ions) m/z 460 (M+H), and


Example 95 title compound (26.7 mg, 36% yield, the other diastereomer) as a colorless solid: LC/MS 100% Area; LC/MS (electrospray, + ions) m/z 460 (M+H).

10

The following compounds were prepared employing procedures as described in Examples 96 to 101.

Example No.	Structure	M+H positive ions
96		405
97		400
98		405
99		391
100		446

101




466

The following compounds were prepared employing the procedures described in Examples 102 to 117 and general procedures detailed in the working examples.

5

Example No.	Structure	M+H positive ions
102	<chem>CC(C)(N)C(=O)N[C@@H](COc1ccccc1)c2nc3c(ncn3N)C(Cc4ccccc4)C#N2</chem>	460
103	<chem>CC(C)(N)C(=O)N[C@@H](COc1ccccc1)c2nc3c(ncn3N)C(Cc4ccccc4)C#N2</chem>	460
104	<chem>CC(C)(N)C(=O)N[C@@H](COc1ccccc1)c2nc3c(ncn3N)C(Cc4ccccc4)C#N2</chem>	448
105	<chem>CC(C)(N)C(=O)N[C@@H](COc1ccccc1)c2nc3c(ncn3N)C(Cc4ccccc4)C#N2</chem>	448


106



Chiral

462


107



Chiral

462


108



Chiral

462


109



Chiral

462


110



Chiral

512


111



Chiral

530


112



Chiral


530

113




416

114




372

115




448

116




448

117




384




Example 118

5

A.




To a solution of crude



- 10 (prepared as described in Example 7 Part C) (7.84 g, 25.37 mmol) in 150 ml of dry ethyl alcohol was added methyl isothiocyanate (2.04g, 27.91 mmol). The reaction was heated to reflux for 6 h. The reaction was allowed to cool to room temperature and the solvent was removed via
- 15 the evaporator. The residue was then treated with 1N NaOH (70 ml), the resulting mixture was heated to reflux (113°C oil bath) for 12 h. The reaction was cooled to room temperature and the solvent was removed via vacuum. The residue was treated with 300 ml of methyl
- 20 chloride/methanol (100:7.5). The mixture was stirred at room temperature for 1 h and filtered. The filtrate was concentrated and co-evaporated with toluene to give crude title compound (7.54g). HPLCb retention time (rt) = 2.37 min. LC/MS (electrospray, + ions) m/z 265 (M+H).


25

B.




To a solution of Part A compound (7.54g, 28.6 mmol) and Boc-methylalanine carboxylic acid (6.96g, 34.3 mmol, Sigma) in 100 ml of DMF was added HOAt (5.05g, 37.1 mmol) and EDAC (8.21g, 42.8 mmol) at room temperature under nitrogen and the reaction mixture was stirred at rt. overnight. The solvent was removed via vacuum and the residue was diluted with water (100 ml). The mixture was extracted with 1/1 ethyl acetate/ether (3x200 ml), the organic layer was dried over magnesium sulfate, and filtered. The filtrate was stored at 0°C and the resulting solid recovered by filtration to give Part B compound (1.45g). The resulting filtrate was then concentrated and the residue was purified by flash chromatography, eluting with hexanes/ethyl acetate (2:1) to give further Part B compound as a white solid (6.92g). HPLCb retention time (rt) = 3.19 min. LC/MS (electrospray, + ions) m/z 450 (M+H).

C.




To a solution of Part B compound (67.4 mg, 0.15 mmol) in 1 ml of dry dioxane was added triethyl amine (30.3 mg, 0.3 mmol) and 2-cyanobenzyl bromide (58.8 mg, 0.3 mmol) at rt. The reaction was shaken overnight and

the solvent was removed via vacuum to give the intermediate:



The residue was then treated with TFA/dichloro-  
5 methane (1 ml, 1:3) at rt for 2h. The solvent was  
evaporated and the reaction mixture was purified by a SCX  
column (2 g) as follows: The column was conditioned by  
rinsing with methanol (10 mL). The reaction mixture in 1  
mL of methanol was loaded onto the column followed by  
10 methanol (30 mL). The product was then eluted with 2 N  
ammonia in methanol (7 mL). The solvent was removed from  
the sample by the use of a speed vacuum to give the title  
compound (53 mg, 76%) as an oil: HPLCb rt = 3.15 min,  
LC/MS (electrospray, + ions) m/z 465 (M+H).  
15


#### Example 119



20 To a solution of Example 118 Part B compound (67.4  
mg, 0.15 mmol) in 1 mL of dry dioxane was added triethyl  
amine (30.3 mg, 0.3 mmol) and 4-(2-bromoisovaleryl)-  
morpholine (75 mg, 0.3 mmol) at rt. The reaction was  
shaken overnight, then potassium carbonate (103 mg, 0.75  
25 mmol) was added and the reaction was continually shaken  
for another 24 h. The solid was filtered off, the

filtrate was collected, and the solvent was removed via vacuum. The residue was then treated with TFA/methylene chloride (1 ml, 1:3) at rt for 2h. The solvent was evaporated and the reaction mixture was purified by a SCX column (2 g) as follows: The column was conditioned by rinsing with methanol (10 mL). The reaction mixture in 1 ml of methanol was loaded onto the column followed by methanol (30 mL). The product was then eluted with 2 N ammonia in methanol (7 mL). The solvent was removed from the sample by the use of a speed vacuum to give the title compound (41 mg, 53%) as an oil: HPLCb rt = 3.06 min, LC/MS (electrospray, + ions) m/z 519 (M+H).

#### Example 120



To a solution of Example 118 Part B compound (67.4 mg, 0.15 mmol) in 1 ml of dry dioxane was added triethyl amine (30.3 mg, 0.3 mmol) and 2-bromobutyrolactone (49.5 mg, 0.3 mmol) at rt. The reaction was shaken overnight, then potassium carbonate (103 mg, 0.75 mmol) was added and the reaction was continually shaken for another 24 h. The solid was filtered off, the filtrate was collected, and the solvent was removed via vacuum. The reaction mixture was purified by a short silica gel cartridge column (2 g) as follows: The column was conditioned by rinsing with ethyl acetate (10 ml). The reaction mixture in 1 ml of ethyl acetate was loaded onto the column followed by ethyl acetate (7 ml). The solution was collected and the solvent was removed from the sample by the use of a speed vacuum. The residue was then treated with TFA/methyl


chloride (1 ml, 1:3) at rt for 2h. The solvent was removed via vacuum to give the title compound as TFA salt (47.1 mg, 57%): HPLC rt = 2.56 min, LC/MS (electrospray, + ions) m/z 434 (M+H).

5

In a manner analogous to that of compounds of Example 118 to 120, compounds of Examples 121 to 172 listed in the table below were prepared from Example 118 Part B compound (0.15 mmol) and the respective bromide (0.3 mmol). Examples 159-172 required further elaboration before deprotection with TFA. A few examples were purified by preparative HPLC, eluting with a gradient system of methanol and water with 0.2% trifluoroacetic acid. Examples 121 to 172 were isolated as trifluoroacetic acid salts or as a free base.

15


Examples 121 to 172



20


Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
121		498
122		508
123		545
124		450

125




408

126




422

127




498

128




436

129




448

130




480

131




450

132




454

133




554

134




461

135
















484


136



544


137		497
138		462
139		675
140		460
141		478
142		492
143		508
144		475
145		490
146		501
147		594
148		571
149		498

150




516

151




465

152




465

153




485

154




470

155




389

156



406

157




463

158




440

159




497

160




539

161




555

162












565

163



536















164		576
165		519
166		535
167		557
168		571
169		606
170		606
171		526
172		547












Compounds of Examples 173 to 216 listed in the table below were prepared in a manner analogous to that of compounds of Example 118 to 172.

5


#### Examples 173 to 216



Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
173		472
174		531
175		562
176		574
177		585
178		556
179		567
180		660
181		651
182		564
183		636


184		582
185		531
186		551
187		551
188		564
189		529
190		536
191		531
192		558
193		531
194		517

195




474

196




560

197




491

198




564

199




473

200




574

201




611

202




516

203




488

204




574

205




620

206




546

207




610

208




563

209




528

210




585

211




741

212




624

213




628

214




526

215




544

216




564


Example 217

5

A.




10 *N*-Methyl morpholine (1.86 mL, 16.9 mmol) and isobutyl chloroformate (2.19 mL, 16.9 mmol) were added to a solution of *N*-Boc-*O*-benzyl-*D*-serine (5 g, 16.9 mmol, ChemImpex) in THF (35 mL), cooled at -18°C under nitrogen. After stirring the mixture for 0.5 h. at -18°C, a solution of *N*-Cbz-1-3-diaminobutane hydrochloride •HCl




15 (4.14 g, 16.9 mmol) and *N*-methyl morpholine (1.86 mL, 16.9 mmol) in DMF (10 mL) was added and the resulting mixture was allowed to warm up to -5°C over a period of 4 h. The mixture was filtered and the solution was evaporated near to dryness. The residue was taken up in EtOAc and washed with 5% NaHCO<sub>3</sub> (3x50 mL), water and brine. The organic solution was dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated to afford Part A compound (7.84 g, 97% yield): LC/MS (electrospray, + ions) m/z 486 (M+H).

B.




Diethylazodicarboxylate (2.57 mL, 16.3 mmol) and  
5 azidotrimethylsilane (2.16 mL, 16.3 mmol) were added to a  
solution of Part A compound (7.92 g, 16.3 mmol) and  
triphenylphosphine (4.28 g, 16.3 mmol) in THF (60 mL),  
previously cooled at 0°C under nitrogen. The mixture was  
stirred at rt overnight and then cooled at 0°C. Another  
10 equivalent each of triphenylphosphine (4.28 g),  
diethylazodicarboxylate (2.57 mL) and azidotrimethylsilane  
(2.16 mL) was added and stirring was continued for an  
additional 24 h at rt. The reaction mixture was cooled at  
0°C and an aqueous solution (75 ml) of ammonium cerium  
15 (IV) nitrate (2.63 g/100 mL) was added and stirred for 1  
h. The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the  
organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and  
chromatographed (SiO<sub>2</sub> 230-400 mesh, 1/1 hexanes/EtOAc) to  
give the impure Part B compound. The contaminated  
20 tetrazole was dissolved in CHCl<sub>3</sub> and ether was added to  
precipitate Part B compound (2.6 g) as a colorless solid:  
MS (electrospray, + ions) m/z 511 (M+H).

C.



To a solution of Part B compound (2.5 g, 4.80 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added a 4M HCl/dioxane solution (14.7 mL) and stirred for 3 h. The solvents were removed at reduce pressure, the residue concentrated from CH<sub>2</sub>Cl<sub>2</sub> and the residue was taken up in minimal *i*-PrOH. Brine (150 mL) was added and the pH of the aqueous solution was adjusted to 10 by addition of 1M K<sub>2</sub>CO<sub>3</sub>. The aqueous solution was extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated to provide the amine intermediate,




used without further purification in the subsequent reaction.


*N*-Boc-methyl alanine (1.49 g, 7.35 mmol, Sigma), EDAC (1.14 g, 7.35 mmol), HOAt (1 g, 7.35 mmol), 1,2-DCE and DMF (1.29 mL) were mixed at 0°C and stirred for 15 min. The resulting solution was transferred to a 0°C solution of the crude amine intermediate (prepared above) in 1,2-DCE and the mixture was stirred overnight at rt. The reaction mixture was diluted with EtOAc and washed with saturated NaHCO<sub>3</sub>, water and brine. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and chromatographed (SiO<sub>2</sub> 230-400 mesh, 1/1 hexanes/EtOAc) to provide Part C compound (1.58 g) as a colorless oil: MS (electrospray, + ions) *m/z* 596 (M+H).



D.




A methanol (3 mL) solution of Part C compound (100  
5 mg, 0.17 mmol) and 5% palladium on carbon (50 mg) was  
hydrogenated for 1 h at 45 psi hydrogen. The reaction  
mixture was filtered through Celite and concentrated



affording intermediate compound  
(68.1 mg). The resulting residue could be purified by  
10 auto Prep HPLC to give pure intermediate compound (47.6  
mg): MS (electrospray, + ions) m/z 462 (M+H).


A CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and DMF (0.5 mL) solution of




hydantoin

(25.3 mg, 0.16 mmol), EDAC (31.3 mg, 0.16 mmol), and  
15 dimethylamino pyridine (DMAP) (20.4 mg, 0.16 mmol) was  
added to a CH<sub>2</sub>Cl<sub>2</sub> (1 mL) solution of the intermediate  
compound (75 mg, 0.16 mmol) at rt under nitrogen. The  
reaction mixture was allowed to stir at rt overnight and  
the volatiles were removed under vacuum. The resulting  
20 residue was dissolved in methanol and passed through a SCX  
resin column to give Part D compound (72 mg): MS  
(electrospray, + ions) m/z 602 (M+H).

E.




A CH<sub>2</sub>Cl<sub>2</sub> solution of Part D compound (64.3 mg,  
 5 0.107 mmol) was treated with an HCl solution (3.0 mL; 10.2  
 ml of AcCl in 40 ml of 3/2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) for 3 h and  
 evaporated in vacuo. The residue was coevaporated four  
 times with CH<sub>2</sub>Cl<sub>2</sub> to give the title compound (51.4 mg) as  
 a white solid: LC/MS (electrospray, + ions) m/z 502  
 10 (M+H).

Example 218

15

A.




N-Methyl morpholine (2.23 mL, 20.3 mmol) and  
 20 isobutyl chloroformate (2.64 mL, 20.3 mmol) were added to  
 a solution of N-Boc-O-benzyl-D-serine (6 g, 20.3 mmol,  
 ChemImpex) in THF (30 mL), cooled at -20°C under nitrogen.  
 After stirring the mixture for 45 min. at -20°C, a  
 solution of ethanolamine (1.3 g, 21.4 mmol) in THF (20 mL)  
 25 was added and the resulting mixture was allowed to warm up  
 to rt over a period of 4 h. After stirring overnight, the  
 mixture was filtered and the solution was evaporated near

to dryness. The residue was purified by flash column chromatography (silica gel), eluting with hexanes:ethyl acetate (1:4) to afford the Part A compound (7.3 g): LC/MS (electrospray, + ions) m/z 339 (M+H).

5


B.



To a CH<sub>2</sub>Cl<sub>2</sub> (20 mL) solution of Part A compound (3.60 g, 10.7 mmol) at rt under nitrogen was added pyridine (3.44 mL, 42.6 mmol) and acetic anhydride (1.1 mL, 11.7 mmol). After 14 h, the volatiles were removed under vacuum. The resulting residue was purified by flash column chromatography (silica gel), eluting with hexanes:ethyl acetate (2:1) to afford the Part B compound (3.85 g): LC/MS (electrospray, + ions) m/z 381 (M+H).

15

C.




20

To a THF (30 mL) solution of Part B compound (3.83 g, 10.1 mmol) was added triphenylphosphine (2.64 g, 10.1 mmol), diethylazodicarboxylate (1.24 mL, 10.1 mmol) and azidotrimethylsilane (1.34 mL, 10.1 mmol). The mixture was stirred at room temperature (rt) for 24 h and then another equivalent each of triphenylphosphine (2.64 g), diethylazodicarboxylate (1.24 mL) and azidotrimethylsilane (1.34 mL) was added and stirring was continued for an additional 24 h at rt. The volatiles were removed under vacuum and the residue was purified by flash chromatography (SiO<sub>2</sub> 230-400 mesh, 8/1 hexanes/EtOAc) to

30


give somewhat impure intermediate compound



(2.71 g). The contaminated tetrazole was sufficiently pure for the subsequent reaction.


- 5 A 4M HCl/dioxane solution (10 mL) was added to the above intermediate compound (2.71 g,  $\leq 6.69$  mmol) and stirred for 1.5 h. The solvents were removed at reduce pressure and the residue co-evaporated twice with a mixture of toluene and methanol to provide Part C compound
- 10 (2.28 g) as a colorless solid: LC/MS (electrospray, + ions) m/z 381 (M+H).

D.



15

- To a CH<sub>2</sub>Cl<sub>2</sub> (25 mL) solution of Part C compound (2.28 g, 6.7 mmol) was added diisopropylethylamine (2.33 mL, 13.4 mmol), *N*-Boc-methyl alanine (2.04 g, 10 mmol, ChemImpex), HOAt (1.36 g, 10 mmol), and EDAC (1.92 g, 10 mmol) and the mixture was stirred overnight at rt. The reaction mixture was quenched with saturated ammonium chloride, and the aqueous layer was washed three times with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were washed with
- 20 brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated to provide
- 25




intermediate compound

sufficiently pure for the subsequent reaction.

The above intermediate compound in a solution of THF/MeOH (20 mL, 20/1) was treated with 2N lithium hydroxide (10 mL) for 1h. After cooling to 0°C, the reaction mixture was neutralized to pH 7 and the aqueous layer extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 times 100 mL). The combined organics were dried (MgSO<sub>4</sub>), filtered, and concentrated under vacuum. The residue was purified by flash chromatography (SiO<sub>2</sub> 230-400 mesh, 1/1 hexanes/EtOAc) to give Part D compound (1.69 g): MS (electrospray, + ions) m/z 449 (M+H).

E.




15

To a THF (20 mL) solution at 0°C of Part D compound (1.34 g, 3.0 mmol) was added pyridine (0.53 mL, 6.6 mmol). After 3 min, a THF (10 mL) solution of para-nitrochloroformate (1.33 g, 6.6 mmol) was added and the mixture was stirred at 0°C for 1 h and 2 h at rt. The reaction mixture was then filtered and the filtrate concentrated under vacuum. The resulting residue was purified by flash chromatography (SiO<sub>2</sub>, 230-400 mesh) eluting with 1/1 hexanes/EtOAc to give Part E compound (1.52 g): LC/MS (electrospray, + ions) m/z 614 (M+H).

20

25


F.



To a THF (20 mL) solution of Part E compound (3.4 g, 5.55 mmol) was added NCCCCCO (0.74 g, 8.32 mmol) and after 3 h at RT, the volatiles were removed under vacuum. The resulting residue was purified by flash chromatography (SiO<sub>2</sub> 230-400 mesh) eluting with 1/2 hexanes/EtOAc (4/1 to 1/1) to give pure Part F compound (2.325 g), as well as some less pure material (1.03 g): LC/MS (electrospray, + ions) m/z 564 (M+H).

10


G.



Part F compound (1.59 g, 2.82 mmol) was treated  
15 with a 4N HCl/dioxane solution (15 mL) for 1.5 h. The  
volatiles were evaporated *in vacuo* to give the title  
compound (1.44 g) as a colorless solid: LC/MS  
(electrospray, + ions) *m/z* 464 (M+H).


The above intermediate was treated with a 4N HCl/dioxane solution for 3 h and evaporated *in vacuo*. The residue was purified by preparative HPLC to give the title compound (85 mg) as a white solid: MS (electrospray, + ions)  $m/z$  566 (M+H).

25           The intermediate Example 218, Part F can be prepared with the modified conditions as described in Example 219.

Example 219


5

A.



To a THF (250 mL) solution of Example 218 Part B compound (47.0 g, 120 mmol) cooled to 1°C was added triphenylphosphine (32.6 g, 123 mmol) and diisopropylethylamine (5.4 mL, 32 mmol). Diethylazodicarboxylate (19.6 mL, 123 mmol, DEAD) was added dropwise over 15 minutes keeping temperature under 5°C followed by the addition azidotrimethylsilane (16.5 mL, 123 mmol). The mixture was stirred at room temperature (rt) for 24 h and then another equivalent each of triphenylphosphine (32.6 g), DEAD (19.6 mL) and azidotrimethylsilane (16.5 mL) was added as before and stirring was continued for an additional 24 h at rt. Another equivalent each of triphenylphosphine (32.6 g), DEAD (19.6 mL) and azidotrimethylsilane (16.5 mL) was added as before and stirring was continued for an additional 24 h at rt. The reaction was cooled to 1°C and quenched with a solution of ammonium cerium(IV) nitrate (136 g) in water (500 mL) over 45 minutes. The resulting mixture was filtered through celite and concentrated in vacuo and extracted twice with ethyl acetate (250 mL). The combined organic extracts were washed with brine, dried over magnesium sulfate and concentrated in vacuo to give an orange oil (236 g). This oil was dissolved in methyl t-butyl ether (1 L, MTBE), washed twice with 1N NaOH (500 mL), once with water (500

mL) and once with brine (500 mL), dried over magnesium sulfate and concentrated *in vacuo* to give a yellow solid (166 g). This solid was purified through a silica gel pad (500 g, 230-400 mesh) eluting with 15-40% ethyl acetate in hexanes to give the acetate intermediate as a yellow oil




(20.6 g, 41% yield).

A 4M HCl/dioxane solution (325 mL) was added to a solution of the above intermediate compound (106.6 g, 263 mmol, combined from numerous reactions) in  $\text{CH}_2\text{Cl}_2$  (600 mL) and stirred overnight. The solvents were removed *in vacuo* and the residue co-evaporated twice with a mixture of toluene and methanol to provide desired compound (89.3 g, 100% yield)) as a brown foam: LC/MS (electrospray, + ions)  $m/z$  381 (M+H).


15

B.



20 To a  $\text{CH}_2\text{Cl}_2$  (1 L) solution of Part A compound (80.0 g, 262 mmol) was added diisopropylethylamine (120 mL, 262 mmol), *N*-Boc-methyl alanine (80.0 g, 393 mmol), HOAt (53.5 g, 393 mmol), and EDAC (75.3 g, 393 mmol) and the mixture was stirred 1 hour at rt. The reaction mixture was washed  
25 with water (500 mL), 0.5 M HCl (500 mL) and saturated  $\text{NaHCO}_3$  (500 mL), dried ( $\text{MgSO}_4$ ), and evaporated to provide intermediate compound as a yellow foam (146.8 g)






sufficiently pure for the subsequent reaction.

The above intermediate compound in a solution of THF/MeOH (1.25 L, 4/1) was placed in a water bath and 2N lithium hydroxide (65 mL) was added over 15 minutes maintaining the internal temperature below 25°C. The reaction stirred at room temperature for 1h. The reaction mixture was concentrated in vacuo to an oil/solid mixture which was dissolved in water (500 mL) and the pH was adjusted to 5.5 with 1N HCl (40 mL). Dichloromethane was added (500 mL) and the mixture stirred for 30 minutes. The organic layer was separated and the aqueous layer extracted with CH<sub>2</sub>Cl<sub>2</sub> (250 mL). The combined organics were washed with water (400 mL) and brine (400 mL), dried (MgSO<sub>4</sub>), filtered, and concentrated under vacuum to give Part D compound (123.2 g) as a light yellow foam: MS (electrospray, + ions) m/z 449 (M+H).

C.




20

To a CH<sub>2</sub>Cl<sub>2</sub> (350 mL) solution at 0°C of Part B compound (123 g, 260 mmol) was added pyridine (32.0 mL, 391 mmol). After 10 min, a CH<sub>2</sub>Cl<sub>2</sub> (150 mL) solution of para-nitrochloroformate (63.0 g, 313 mmol) was added over 1 hour, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was washed with water (300 mL), 1N HCl (300 mL), and brine (300 mL), dried

(MgSO<sub>4</sub>), and concentrated *in vacuo* to a brown oil (187.3 g). This crude material was purified through a silica gel pad (500 g, 230-400 mesh) eluted with 30-50% ethyl acetate in hexanes to give Part E compound as an off-white foam (130.98 g, 81.9% yield): LC/MS (electrospray, + ions) m/z 614 (M+H).

D.











10
















To a THF (600 mL) solution of Part C compound (117.5 g, 190 mmol) in a water bath was added aminobutan-1-ol (19.4 mL, 210 mmol) dropwise over 30 minutes and after 1 h at RT, an additional aliquot of 4-amino-1-butanol (1.7 mL, 18 mmol) was added. After stirring an additional 30 minutes at rt, the volatiles were removed under vacuum. The resulting residue was dissolved in EtOAc (600 mL) and washed with 1N HCl (200 mL), saturated NaHCO<sub>3</sub> (1 x 400 mL and 2 x 200 mL, emulsion), and brine (3 x 200 mL, slow separation), dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give an oil. This oil was dissolved in MTBE (600 mL) and washed with saturated NaHCO<sub>3</sub> (1 x 300 mL), and brine (2 x 500 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated *in vacuo* to give a yellow oil (156.52 g). This oil was dissolved in MTBE (1 L) and washed with 1N NaOH (500 mL) and water. The organic layer was washed again with 1N NaOH (500 mL) and the combined NaOH washes were extracted with MTBE (400 mL). The combined MTBE layers were washed with water (500 mL) and brine (500 mL), dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give pure title Part D compound as a light yellow oil (103.08 g, 95.5% yield): LC/MS (electrospray, + ions) m/z 564 (M+H).















Examples 220-352 were prepared in a manner analogous to that of compounds described previously in the invention and by methods known in the art.

5
















Examples 220 to 352


















Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
220		466
221		478
222		505
223		505
224		418
225		415
226		434
227		433

228		482
229		476
230		475
231		590
232		474
233		544
234		518
235		462
236		442
237		447
238		434
239		464
240		448
241		502
242		479

















243		511
244		574
245		485
246		462
247		463
248		349
249		524
250		496
251		448
252		434
253		462
254		498
255		496
256		466



















257		498
258		497
259		542
260		512
261		512
262		511
263		462
264		511
265		547
266		404
267		390
268		418
269		518
270		510
271		496














272		510
273		391
274		524
275		489
276		543
277		492
278		506
279		513
280		527
281		582
282		553
283		520
284		476
285		448
286		462

287		476
288		450
289		464
290		480
291		478
292		432
293		510
294		510
295		524
296		562
297		562
298		630
299		512
300		475
301		524
302		524
303		462




304		524
305		462
306		573
307		552
308		478
309		492
310		506
311		520
312		482
313		524
314		504
315		539
316		553
317		532
318		518
319		518

320		554
321		468
322		553
323		526
324		386
325		372
326		567
327		508
328		478
329		474
330		553
331		440
332		498
333		540
334		537
335		528
336		450
337		462

338		480
339		536
340		540
341		563
342		507
343		556
344		494
345		549
346		542
347		531
348		549
349		377
350		439


Example No.	Structure	M+H positive ions
-------------	-----------	-------------------

351




508

352




477

Example 353


5

A.



To a solution of Compound 218 Part F (929mg, 1.65mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (8mL) was added pyridine (400μl, 4.95mmol) and DMAP (40mg, 0.33mmol) followed by the addition of benzoyl chloride (287μl, 2.48mmol). After stirring for 5h at room temperature, the solvent was removed *in vacuo*, and the residue was purified by flash column chromatography (SiO<sub>2</sub>, 20%-50% EtOAc in hexanes) to give the desired product (1.09g, 99% yield).


B.




- 5 To a solution of Part A compound (580mg) in CH<sub>2</sub>Cl<sub>2</sub> (3ml) was added 4N HCl in dioxane (3ml). The reaction mixture stirred at room temperature for 1.5h. The solvent was removed *in vacuo* to give the desired product. (469mg, 89% yield). LC/MS (electrospray, + ions) m/z 567 (M+H).

10

Example 353, alternate preparation



A.




15

- To a solution of Example 219 compound (102.66 g, 182 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (650mL) was added pyridine (37.0 mL, 455 mmol) and DMAP (2.23g, 18.2 mmol). The flask was immersed in a water bath and benzoyl chloride (25 mL, 219 mmol) was added dropwise over 20 minutes. After stirring for 2h at room temperature, additional benzoyl chloride (6.0 mL, 54mmol) was added dropwise and the reaction continued to stir at room temperature overnight. The reaction mixture was washed with 1N HCl (300 mL), saturated NaHCO<sub>3</sub> (300 mL), and brine (300 mL), dried (MgSO<sub>4</sub>), and concentrated *in vacuo*, and the residue


was purified by silica gel pad filtration ( $\text{SiO}_2$ , 650g, 10%-85% EtOAc in hexanes) to give the desired product (116.34 g, 97% yield) as a colorless foam.

5 B.




To a solution of Part A compound (116.0 g, 174 mmol) in  $\text{CH}_2\text{Cl}_2$  (700 ml) was added 4N HCl in dioxane (218 ml, 870 mmol) dropwise over 30 minutes. The reaction mixture stirred at room temperature for 4 h. The solvent was removed *in vacuo*. The residue was dissolved in water (500 mL) and washed with MTBE (250 mL). The aqueous layer was mixed with  $\text{CH}_2\text{Cl}_2$  (300 mL) and the pH adjusted to 11 with 1N NaOH (190 mL). The  $\text{CH}_2\text{Cl}_2$  layer was separated and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (300 mL). The combined  $\text{CH}_2\text{Cl}_2$  layers were washed with brine (2 x 300 mL), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo* to give the desired product (92.8 g, 94% yield) as a yellow oil. LC/MS (electrospray, + ions) m/z 567 (M+H).

#### Example 354




25

A.



- 5 To a solution of Compound 219 (100 mg, 0.18 mmol) in 1,2-dichloroethane (250μl) at 0°C is added a solution of 2-phenylpropionic acid (50μl, 0.37 mmol), EDAC (69mg, 0.36 mmol), and DMAP (44mg, 0.36 mmol) in 1,2-dichloroethane (250μl). The reaction mixture stirred for 36 hours
- 10 at room temperature. The reaction was diluted with EtOAc (50 ml) and washed with saturated NaHCO<sub>3</sub> (2x25 ml), water (25 ml) and brine (25 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give the crude product (128 mg). This material was purified by flash column chromatography eluted with 1:1
- 15 hexanes:EtOAc to give the product (122.4 mg, 99% yield).


B.



- 20 Part A compound (122 mg, 0.18 mmol) was stirred in a 10%TFA in CH<sub>2</sub>Cl<sub>2</sub> solution (4.0 ml) for 2.5h. The solvent was removed *in vacuo*. The residual TFA was removed by coevapoation with CH<sub>2</sub>Cl<sub>2</sub> (4x3 ml) and MeOH (2x3
- 25 ml). The residue was dissolved in MeOH and heated at 60°C for 8h. The solvent was removed *in vacuo*, and the residue was purified by preparative HPLC (20%-100% B, 12min. gradient, 20ml/min, YMCS5 ODS 20x100mm) to give the desired product (115.4 mg, 93% yield). LC/MS
- 30 (electrospray, + ions) m/z 596 (M+H).

In a manner analogous to that of compounds of Example 353 and 354, compounds of Examples 355 to 392 listed in the table below were prepared from Example 218 Part F compound and the respective acid or acid chloride.

5















Examples 355 to 392















Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
355		534
356		548
357		582
358		610
359	$\text{X}_1\text{-CH}_3$	506
360		548
361		520
362		586
363		604
364		636




WO 00/54729

PCT/US00/05704







365		582
366		596
367		582
368		596
369		644
370		598
371		626
372		603
373		548
374		562
375		582
376		588
377		644
378		618


379		618
380		660
381		660
382		574
383		560
384		546
385		594
386		574
387		560
388		636
389		652
390		650
391		534
392		546











Compounds of Examples 393 to 424 listed in the tables below were prepared in a manner analogous to that of compounds of Example 353 and 354.

Examples 393 to 424

Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
393		566
394		580
395		572
396		580
397		594
398		560
399		596
400		606
401		610
402		610
403		586
404		598

405		648
406		648
407		546
408		700
409		580
410		544



Example No.	X <sub>1</sub> -R	M+H positive ions
411		568
412		596
413		582
414		630
415		625
416		597
417		644
418		665
419		660
420		610

421




650

422



648

423




624

424




636


Example 425

5

A.



To a solution of methyl R-(+)-3-BOC-2,2-dimethyl-4-




10 oxazolidinone carboxylate

(2.05 g, 7.75 mmol, Aldrich, 98%) in THF (24.0 mL) was added a light suspension of calcium chloride (437 mg, 3.94 mmol) in EtOH (16.5 mL) and the resulting solution was cooled at 0°C. Sodium borohydride was added and the mixture was stirred at 0°C for 2h. and for 4h at rt. The final mixture was cooled at 0°C and potassium phosphate buffer (pH 3, 40 mL) was added. The aqueous mixture was stirred for 30 min at rt. and then extracted with CH<sub>2</sub>Cl<sub>2</sub>

(3x70 mL). The combined organic phase was dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to a crude which was chromatographed ( $\text{SiO}_2$  230-400 mesh, 3/2 to 1/1 hexanes/ $\text{EtOAc}$ ) to give the starting ester (409 mg, 20% recovery) and the desired  
 5 alcohol (1.46 g, 80% yield):  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm, rotamers) 4.18-3.56 (4m, 6H), 1.55 and 1.50 (2s, 15 H).

B.




10

To a solution of Part A compound (2.90 g, 12.5 mmol), 2-hydroxypyridine (1.54 g, 15.7 mmol) and triphenylphosphine (4.11 g, 15.7 mmol) in THF (27 mL) was added  
 15 diethylazo-dicarboxylate (2.47 mL, 15.7 mmol) dropwise. The solution was stirred at rt. for 13.5 h. and then partially evaporated. The remaining solution was passed through a  $\text{SiO}_2$  (230-400 mesh) column, eluting with 7/3 hexanes/ $\text{EtOAc}$ , to provide the desired compound (1.88 g,  
 20 49% yield) as a solid:  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm, rotamers) 8.15 (m, 1H), 7.57 (m, 1H), 6.88 (m, 1H), 6.74 (d,  $J = 8.2$  Hz, 1H), 4.50 (m, 1H), 4.25-3.95 (several m, 4H), 1.61-1.45 (several s, 15H).


25

C.




A solution of Part B compound (434 mg, 1.41 mmol) and p-toluenesulfonic acid monohydrate (295 mg, 1.55 mmol) in dry MeOH (14.1 mL) was heated at 35°C for 7 h and then stirred at rt. for 3.8 h. After cooling at 0°C, a 1M K<sub>2</sub>CO<sub>3</sub> solution (0.8 mL) was added and the mixture volume was reduced in vacuo to 4 mL. Brine (50 mL) was added and the pH of the aqueous solution was adjusted to 10 by addition of 1M K<sub>2</sub>CO<sub>3</sub>. Extraction with CH<sub>2</sub>Cl<sub>2</sub> (4x50 mL), drying (Na<sub>2</sub>SO<sub>4</sub>), evaporation and chromatography (SiO<sub>2</sub>, 230-400 mesh, 1/1 hexanes/EtOAc) of the crude gave starting Part B compound (93 mg, 21% recovery) and the desired alcohol (249 mg, 66% yield) as a colorless solid: LC-MS 99% Area; LC/MS (electrospray, + ions) m/z 269 (M+H).

D.




To a solution of Part C compound (1.34 g, 5.00 mmol) and triphenylphosphine (1.58 g, 6.02 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (34.0 mL), cooled at 0°C, was added diethylazodicarboxylate (0.95 mL, 6.03) dropwise and the mixture was stirred at rt. for 3.75 h. The mixture volume was reduced to 8 mL in vacuo and the remaining solution was passed through a SiO<sub>2</sub> (230-400 mesh) column, eluting with 7/3 hexanes/EtOAc to obtain the desired compound (862 mg, 70% yield) as a yellow oil: <sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>, ppm) 8.13 (m, 1H), 7.57 (m, 1H), 6.88 (m, 1H), 6.78 (d, *J* = 8.2 Hz, 1H), 4.45 (dd, *J* = 11.6, 4.4 Hz, 1H), 4.36 (dd, *J* = 11.6, 5.5 Hz, 1H), 2.88 (m, 1H), 2.35 (d, *J* = 6.0 Hz, 1H), 2.22 (d, *J* = 3.9 Hz, 1H), 1.43 (s, 9H).

E.



- 5 To a suspension of Part D compound (97 mg, 0.39 mmol) and potassium cyanide (51 mg, 0.78 mmol) in DMSO (2.6 mL) was added water (7  $\mu$ L) and the mixture was heated at 30-40 °C for 11 h.. Stirring was continued for an additional 9 h. at rt. and the mixture was diluted with 9/1
- 10 hexanes/EtOAc (60 mL). The solution was washed with water (2x30 mL) and brine (30 mL), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to give 92 mg crude. Chromatography ( $\text{SiO}_2$  230-400 mesh, 7/3 hexanes/EtOAc) provided starting aziridine (8.0 mg, 8% recovery) and the desired compound (75.6 mg, 72% yield) as
- 15 a colorless solid: LC-MS 100% Area; LC/MS (electrospray, + ions) m/z 278 (M+H).

F.




20

- 15% TFA solution (6.0 mL) was added to a flask containing Part E compound (199 mg, 0.72 mmol) and thioanisole (254  $\mu$ L, 2.16 mmol), and the resulting solution was allowed to stand at rt. for 4 h. The
- 25 solution was evaporated and the residual TFA was removed




by coevaporation with  $\text{CH}_2\text{Cl}_2$  (2x5 mL) and MeOH (3x5 mL), and drying under high vacuum for 1h. The crude amine salt



material, was used without further purification in the subsequent reaction.

- 5        *N*-Boc-*O*-benzyl-D-serine (298 mg, 1.00 mmol), EDAC (194 mg, 1.01 mmol), HOAT (137 mg, 1.01 mmol), 1,2-DCE (0.47 mL), and DMF (940  $\mu\text{L}$ ) were mixed at 0°C and stirred for 15 min. The resulting cloudy solution was transferred, via syringe, to a 0°C solution of the crude
- 10    amine (from above) in 1,2-DCE (400  $\mu\text{L}$ ) and DMF (470  $\mu\text{L}$ ), and the mixture was stirred for 10 min. at 0°C. Diisopropylethylamine was added and stirring was continued at rt. for 36 h. The reaction mixture was diluted with EtOAc (80 mL) and washed with saturated  $\text{NaHCO}_3$  (2x40 mL),
- 15    water (40 mL) and brine (40 mL). The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), evaporated and chromatographed ( $\text{SiO}_2$  230-400 mesh, 1/1 hexanes/EtOAc) to provide the desired compound (311 mg, 95% yield) as a colorless oil:  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm) 8.17 (s, 1H), 7.84 (s, 1H), 7.61 (m, 1H), 7.30 (m, 5H), 6.94 (m, 1H), 6.76 (d,  $J = 7.9$  Hz, 1H), 5.37 (s, 1H), 4.52 (m, 5H), 4.31 (broad s, 1H), 3.88 (m, 1H), 3.58 (m, 1H), 2.80 (m, 2H), 1.43 (s, 9H).
- 20


G.



25

Diethylazodicarboxylate (131  $\mu$ L, 0.83 mmol, DEAD) and  
azidotrimethylsilane (110  $\mu$ L, 0.83 mmol) were added to a  
solution of Part F compound (311 mg, 0.69 mmol) and  
5 triphenylphosphine (217 mg, 0.83 mmol) in  $\text{CH}_2\text{Cl}_2$  (1.4 mL).  
The mixture was stirred at rt. for 22 h. and additional  
amounts of triphenylphosphine (217 mg), DEAD (131  $\mu$ L) and  
azidotrimethylsilane (110  $\mu$ L) were added. Stirring was  
continued for an additional 24 h at rt. and the final  
10 mixture was passed through a  $\text{SiO}_2$  (230-400 mesh) column,  
eluting with 3/2 to 1/1 hexanes/EtOAc, to give the desired  
compound (248 mg, 76%) as a thick colorless oil:  $^1\text{H}$  NMR  $\delta$   
( $\text{CDCl}_3$ , ppm) 8.11 (m, 1H), 7.55 (m, 1H), 7.33 (m, 3H), 7.20  
(m, 2H), 6.92 (t,  $J = 7.9$  Hz, 1H), 6.76 (d,  $J = 8.3$  Hz,  
15 1H), 5.45 (d,  $J = 8.3$  Hz, 1H), 5.44 (m, 2H), 4.73 (dd,  $J =$   
11.6, 4.9 Hz, 1H), 4.63 (dd,  $J = 11.6$ , 8.3 Hz, 1H), 4.50  
(d,  $J = 11.0$  Hz, 1H), 4.44 (d,  $J = 11.0$  Hz, 1H), 4.00 (dd,  
 $J = 8.3$ , 5.5 Hz, 1H), 3.58 (t,  $J = 8.8$  Hz, 1H), 3.04 (dd,  
 $J = 17.0$ , 7.1 Hz, 1H), 2.86 (dd,  $J = 17.0$ , 7.1 Hz, 1H),  
20 1.43 (s, 9H).


H.




25

5% TFA solution (6.2 mL) was added to a flask  
containing the Part G compound (227 mg, 0.47 mmol) and


thioanisole (166  $\mu$ L, 1.41 mmol), and the resulting solution was allowed to stand at rt. for 4 h. The solution was evaporated and the residual TFA was removed by coevaporation with  $\text{CH}_2\text{Cl}_2$  (2x8 mL) and MeOH (3x8 mL), and drying under high vacuum for 1h. The residue was heated at 60°C in MeOH (25 mL) and the solution



concentrated. The crude amine material, used without further purification in the subsequent reaction.

- 10 *N*-Boc-methyl alanine  (125 mg, 0.62 mmol), EDAC (117 mg, 0.61 mmol), HOAT (84 mg, 0.62 mmol), 1,2-DCE (350  $\mu$ L) and DMF (700  $\mu$ L) were mixed at 0°C and stirred for 15 min. The resulting solution was transferred, via syringe, to a 0°C solution of the crude amine (from above) in 1,2-DCE (300  $\mu$ L) and DMF (400  $\mu$ L), and the mixture was stirred at 0°C for 15 min. Diisopropylethylamine (205  $\mu$ L, 1.18 mmol) was added and stirring was continued for 25 h at rt. The final mixture was diluted with EtOAc (75 mL) and washed with saturated  $\text{NaHCO}_3$  (2x30 mL), water (30 mL) and brine (30 mL). The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), evaporated and chromatographed ( $\text{SiO}_2$  230-400 mesh, 3/2 to 2/3 hexanes/EtOAc) to provide desired compound (175.4 mg, 66% yield) as a colorless foam:  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm) 8.13 (m, 1H), 7.57 (m, 1H), 7.42 (d,  $J$  = 8.8 Hz, 1H), 7.31 (m, 3H), 7.19 (m, 2H), 6.93 (m, 1H), 6.77 (d,  $J$  = 8.2 Hz, 1H), 5.75 (m, 1H), 5.49 (m, 1H), 4.97 (s, 1H), 4.73 (dd,  $J$  = 11.6, 5.5 Hz, 1H), 4.66 (dd,  $J$  = 11.6, 7.7 Hz, 1H), 4.50 (d,  $J$  = 11.6 Hz, 1H), 4.43 (d,  $J$  = 11.5 Hz, 1H), 4.04 (m, 1H), 3.52 (t,  $J$  = 8.8 Hz, 1H), 3.02 (dd,  $J$  = 17.1, 6.6 Hz, 1H), 2.85 (dd,  $J$  = 17.0, 7.7 Hz, 1H), 1.50 (s, 3H), 1.44 (s, 3H), 1.36 (s, 9H).

I.




5 Part H compound (492 mg, 0.87 mmol) was treated with 15% TFA (14.0 mL) for 2.5 h and the solution concentrated in vacuo. The residual TFA was coevaporated with CH<sub>2</sub>Cl<sub>2</sub> (3x10 mL) and MeOH (2x10 mL), and drying under high vacuum for 1h.. The residue was heated at 60°C in MeOH (40 mL)

10 for 10 h. and the solution concentrated to give the crude desired compound. Successive precipitation from MeOH/ether provided the desired compound (418 mg). The mother liquor was evaporated and the residue was purified by preparative HPLC (Shimadzu, 10-100% B [MeOH:H<sub>2</sub>O:0.1%



















15 TFA], 30 min. gradient, 20 mL/min. flow rate, 220nm, YMC S5 ODS 20x100mm) to give an additional amount of the title compound (67.6 mg): Combined yield, 96%; LC-MS 99% Area; LC/MS (electrospray, + ions) m/z 465 (M+H).

20 The following compounds were prepared employing the procedures described above and the working Examples.


Examples 426 to 477




25

Example No.	$X_1-R$	M+H positive ions
426		540
427		490
428		462
429		504
430		449
431		450
432		480
433		477
434		463
435		523
436		500
437		466
438		575
439		504
440		464
441		581
442		490
443		478


444




445




446




447




448




449




450




451




452




453




454




455























456











457

















458		476
459		462
460		503
461		546
462		546
463		567
464		601
465		464
466		560
467		517
468		593
469		595
470		518
471		565
472		476

473		552
474		622
475		565
476		579
477		607


Examples 478 to 498

Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
478		566
479		566
480		552
481		519
482		532
483		572
484		510



























485		523
486		574
487		573
488		460
489		516
490		545
491		488
492		501
493		474
494		523
495		574
496		473
497		544
498		508


Examples 499 to 530








Example No.	X <sub>1</sub> -R	M+H positive ions
499		464
500		525
501		496
502		532
503		511
504		496
505		532
506		511

507		544
508		483
509		455
510		482
511		524
512		496
513		565
514		450
515		482
516		497
517		518
518		497
519		539
520		510
521		546


522		525
523		482
524		496
525		496
526		482
527		512
528		512
529		511
530		525

Examples 531 to 554

Example No.	<u>X<sub>1</sub>-R</u>	M+H positive ions
531		581
532		450
533		464
534		497
535		511


536		462
537		476
538		490
539		405
540		344
541		464
542		536
543		536
544		557
545		557
546		506
547		589
548		370
549		521
550		583
551		465
552		426

553




535


554



480


Examples 555 to 583**Example No.****X<sub>1</sub>-R****M+H positive ions**

555




453

556




489

557




511

558




525

559











483

560




465

Example No.	Structure	M+H positive ions
561		481
562		416
563		500
564		498
565		618
566		602
567		601
568		615

- 164 -




580



517


581



Chiral

531


582



Chiral

377

583




Chiral

463

What is claimed is:

1. A compound of the structure



5

including pharmaceutically acceptable salts thereof, prodrug esters thereof, and all stereoisomers thereof,

wherein R<sub>1</sub> is alkyl, aryl, alkenyl, alkynyl,

arylalkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl,

10 arylalkyloxyalkyl, aryloxyalkyl, heteroaryl,

cycloalkylalkoxyalkyl, heteroarylalkyl, cycloheteroalkyl

or cycloheteroalkylalkyl, and these groups may be


optionally substituted by 1,2 or 3-substituents selected from halogen, -OR<sub>8</sub>, -OC(O)R<sub>8</sub>, alkyl, phenyl, phenoxy,

15 halophenyl, -CF<sub>3</sub>, -OCF<sub>3</sub>, -N(R<sub>8a</sub>)C(O)(R<sub>8</sub>), or


-N(R<sub>8</sub>)(R<sub>8a</sub>);

R<sub>1a</sub> is H, alkyl, or cycloalkyl;

X<sub>a</sub> is heteroaryl, which is




20



- A is oxygen, sulfur, -NH-, -N-R<sub>5</sub>, or -NC(O)-R<sub>2</sub>;  
 B is -CR<sub>5b</sub> or -N-;
- 5 Z is a bond or -S-;  
 G is oxygen or sulfur;  
 U is oxygen, sulfur, -NH-, or -N-R<sub>5b</sub>;
- R<sub>2</sub> is alkyl, aryl, alkenyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, heteroaryl, or heteroarylalkyl, and these groups may optionally be substituted by 1,2 or 3-substituents selected from halogen, -OR<sub>8b</sub>, -OC(O)R<sub>8b</sub>, alkyl, phenyl, phenoxy, halophenyl, -CF<sub>3</sub>, -OCF<sub>3</sub>, -N(R<sub>8c</sub>)C(O)(R<sub>8b</sub>), or -N(R<sub>8c</sub>)(R<sub>8b</sub>);
- 10 R<sub>3</sub> is H, halogen, alkyl, aryl, alkenyl, alkynyl, alkaryl, alkoxy, aryloxy or J<sub>1</sub>, and where alkyl, aryl, alkenyl, alkynyl, arylalkyl, alkoxy, or aryloxy may be optionally substituted with 1 to 3 J<sub>1</sub>;
- R<sub>4</sub> and R<sub>4a</sub> are the same or different and are
- 20 independently selected from H, halogen, -CF<sub>3</sub>, alkyl, or aryl;
- R<sub>5</sub> is H, alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, cycloalkylalkoxyalkyl, -SO<sub>2</sub>T<sub>1</sub>, -SO<sub>2</sub>N(T<sub>1a</sub>)T<sub>1</sub>, or
- 25 heteroarylalkyl, and where alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, heteroaryloxyalkyl, cycloalkylalkoxyalkyl, or heteroarylalkyl may be
- 30 independently optionally substituted with 1 to 3 J<sub>1</sub>;
- R<sub>5a</sub> and R<sub>5b</sub> are the same or different and are independently selected from H, alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl,

cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, cycloalkylalkoxyalkyl, heteroarylalkyl, or J1, and where alkyl, aryl, alkenyl, arylalkenyl, alkynyl, arylalkyl, arylalkynyl, cycloalkylalkyl, alkoxyalkyl, aryloxyalkyl, arylalkyloxyalkyl, heteroaryl, heteroaryloxyalkyl, cycloalkylalkoxyalkyl, or heteroarylalkyl may be independently optionally substituted with 1 to 3 J1;

Y is



where x and y are independently 0 to 3 and z is 1 to 3;

$X_c$  is a bond,  $\text{---N-R}_{6a}$  or  $\text{---O-}$ ;

$R_7$  and  $R_{7a}$  are the same or different and are


15 independently selected from H, alkyl,  $\text{---CF}_3$ , phenyl, aryl, arylalkyl, and cycloalkyl; or one or both of  $R_7$  and  $R_{7a}$  can be independently joined to one or both of  $R_9$  and  $R_{10}$  groups (of  $X_b$ ) to form an alkylene bridge of 1 to 5 carbon atoms; or  $R_7$  and  $R_{7a}$  are joined together to form a ring of

20 from 3-7 carbon atoms;

$R_6$ ,  $R_{6a}$ ,  $R_{6b}$ ,  $R_{6c}$ ,  $R_8$ ,  $R_{8a}$ ,  $R_{8b}$ ,  $R_{8c}$ ,  $R_{8d}$ ,  $R_{8e}$ ,  $R_{8f}$ ,  $R_{8g}$ ,  $R_{8h}$ ,  $R_{8i}$ ,  $R_{8k}$ ,  $R_{8l}$ , and  $R_{8m}$  are the same or different and are independently H, alkyl, cycloalkyl, alkenyl or aryl;

25  $R_{8j}$  is H, alkyl, aryl, hydroxy or  $\text{---OC(O)R}_{8k}$ ;

$X_b$  is




$R_9$  and  $R_{10}$  are the same or different and are

30 independently selected from H, alkyl, and substituted alkyl where the substituents may be 1 to 3 hydroxy, 1 to 3


- C<sub>1</sub>-C<sub>10</sub>-alkanoyloxy, 1 to 3 C<sub>1</sub>-6 alkoxy, phenyl, phenoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl; or R<sub>9</sub> and R<sub>10</sub> can together form -(CH<sub>2</sub>)<sub>t</sub>X<sub>d</sub>(CH<sub>2</sub>)<sub>u</sub> - where X<sub>d</sub> is C(R<sub>8h</sub>)(R<sub>8j</sub>), -O- or -N(R<sub>6b</sub>), t and u are independently 1-3;
- 5 R<sub>11</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, -CF<sub>3</sub>, arylalkyl, or aryl, and with the alkyl and aryl groups being optionally substituted with 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-<sub>10</sub>alkanoyloxy, 1 to 3 C<sub>1</sub>-6 alkoxy, phenyl, phenoxy or C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl;
- 10 R<sub>12</sub> and R<sub>13</sub> are independently H, C<sub>1</sub>-C<sub>6</sub>alkyl, -CF<sub>3</sub>, aryl, or halogen, and with the alkyl and aryl groups being optionally substituted with 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-C<sub>10</sub>-alkanoyloxy, 1 to 3 C<sub>1</sub>-6 alkoxy, or C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl;
- J<sub>1</sub> is nitro, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>CN,
- 15 -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)C(O)N(T<sub>1b</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)SO<sub>2</sub>T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OC(O)OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OC(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OC(O)N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>N(T<sub>1a</sub>)SO<sub>2</sub>N(T<sub>1b</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>OT<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>SO<sub>2</sub>T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>SO<sub>2</sub>N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>C(O)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>CH(OH)T<sub>1</sub>,
- 20 cycloheteroalkyl or heteroaryl, with v being 0-5;
- T<sub>1</sub>, T<sub>1a</sub> and T<sub>1b</sub> are the same or different and are independently H, alkyl, alkenyl, alkynyl, lower alkythioalkyl, alkoxyalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, or cycloalkyl, each of
- 25 which may be optionally substituted with 1, 2 or 3 substituents selected from halogen, hydroxyl, -NR<sub>8f</sub>C(O)NR<sub>8g</sub>R<sub>8i</sub>, -C(O)NR<sub>8f</sub>R<sub>8g</sub>, -NR<sub>8f</sub>C(O)R<sub>8g</sub>, -CN, -N(R<sub>8f</sub>)SO<sub>2</sub>R<sub>14</sub>, -OC(O)R<sub>8f</sub>, -SO<sub>2</sub>NR<sub>8f</sub>R<sub>8g</sub>, -SOR<sub>14</sub>, -SO<sub>2</sub>R<sub>14</sub>, alkoxy, -COOH, cycloheteroalkyl, or -C(O)OR<sub>14</sub>; or
- 30 T<sub>1</sub> and T<sub>1a</sub> or T<sub>1</sub> and T<sub>1b</sub> can together form -(CH<sub>2</sub>)<sub>w</sub>X<sub>e</sub>(CH<sub>2</sub>)<sub>z</sub>- where X<sub>e</sub> is -C(R<sub>8m</sub>)(R<sub>8l</sub>), -O-, -S-, -SO-, -SO<sub>2</sub>-, -NC(O)OR<sub>14a</sub>, -NC(O)NR<sub>14a</sub>R<sub>14b</sub>, -NC(O)R<sub>14a</sub> or -N(R<sub>6c</sub>) where w and z are each independently 1-3; with the proviso that T<sub>1</sub> can not be hydrogen when it is connected to
- 35 carbonyl or sulfur, as in -C(O)T<sub>1</sub> or -SO<sub>2</sub>T<sub>1</sub>;

R<sub>14</sub>, R<sub>14a</sub>, and R<sub>14b</sub> are independently C<sub>1</sub>-C<sub>6</sub>alkyl, heteroaryl, or aryl, each optionally substituted with -(CH<sub>2</sub>)<sub>s</sub>OH, with s being 0-5;



with the proviso that where X<sub>a</sub> is


- 5 (1) where one or both of R<sub>7</sub> and R<sub>7a</sub>, and one or both of R<sub>9</sub> and R<sub>10</sub> form an alkylene bridge then where R<sub>5</sub> is -(CH<sub>2</sub>)C(O)N(T<sub>1a</sub>)T<sub>1</sub>, then at least one of T<sub>1a</sub> and T ≠ H; or
- (2) where R<sub>1</sub> is arylalkyl and R<sub>1a</sub> is H and R<sub>5</sub> is
- 10 -(CH<sub>2</sub>)C(O)N(T<sub>1a</sub>)T<sub>1</sub>, then T<sub>1a</sub> or T<sub>1</sub> is other than



- (3) where R<sub>1</sub> and R<sub>7</sub> are each 2-naphthyl-CH<sub>2</sub>-, then R<sub>5</sub> ≠ phenethyl.


15

2. The compound as defined in Claim 1 wherein X<sub>a</sub> is




20

3. The compound as defined in Claim 1 having the structure



4. The compound as defined in Claim 1 having the structure



5


5. The compound as defined in Claim 1 wherein R<sub>1</sub> is aralkyl, arylalkyloxyalkyl, cycloheteroalkylalkyl, aryloxyalkyl or heteroarylalkyl.

10 6. The compound as defined in Claim 1 wherein R<sub>1a</sub> is H or alkyl.

7. The compound as defined in Claim 1 wherein R<sub>1</sub> is arylalkyloxyalkyl and R<sub>1a</sub> is H.


15 8. The compound as defined in Claim 1 wherein R<sub>6</sub> is H.

9. The compound as defined in Claim 7 wherein (1) X<sub>a</sub> is




where R<sub>5</sub> is alkyl or alkenyl or heteroaryloxy-alkyl, each substituted with J<sub>1</sub>, and J<sub>1</sub> is

20 -(CH<sub>2</sub>)<sub>v</sub>OC(O)N(T<sub>1a</sub>)T<sub>1</sub>, -(CH<sub>2</sub>)<sub>v</sub>CN, or heteroaryl; or




(2)  $X_a$  is where A is NH,  $R_4$  and  $R_{4a}$  are H and  $R_3$  is J1; or




(3)  $X_a$  is where B is -N- or  $CR_{5b}$ ; Z is a bond or -S-;  $R_5$  is alkyl optionally substituted with J1; and  $R_{5a}$  is H, or alkyl or aralkyl substituted with 1 to 3 J1;  
 $R_6$  is H.

10. The compound as defined in Claim 1 wherein Y



is where x and y are 0,  $X_c$  is a bond, and  $R_7$  and  $R_{7a}$  are independently alkyl.

11. The compound as defined in Claim 1 wherein  $X_b$  is



15  $R_9$  and  $R_{10}$  are the same or different and are independently selected from H and substituted alkyl where the substituents may be 1 to 2 hydroxyls.

20 12. The compound as defined in Claim 1 wherein J1 is  $-(CH_2)_vCN$ ,  $-(CH_2)_vC(O)N(T_{1a})T_1$ ,  $-(CH_2)_vN(T_{1a})C(O)T_1$ ,  $-(CH_2)_vOC(O)N(T_{1a})T_1$ ,  $-(CH_2)_vN(T_{1a})C(O)N(T_{1b})T_1$ , or heteroaryl, with v being 0-4;

25  $T_1$ ,  $T_{1a}$  and  $T_{1b}$  are the same or different and are independently selected from alkyl, lower alkythioalkyl, alkoxyalkyl, aryl, arylalkyl, heteroarylalkyl, or














cycloheteroalkyl, each of which may be optionally substituted with  $-\text{OC}(\text{O})\text{R}_{8f}$ ,  $-\text{C}(\text{O})\text{NR}_{8f}\text{R}_{8g}$ ,  $-(\text{CH}_2)_s\text{OH}$ , with  $s$  being 0-2,  $-\text{SO}_2\text{NR}_{8f}\text{R}_{8g}$ , or  $-\text{SO}_2\text{R}_{14}$ ; or  $\text{T}_1$  and  $\text{T}_{1a}$  or  $\text{T}_1$  and  $\text{T}_{1b}$  can together form  $-(\text{CH}_2)_w\text{X}_e(\text{CH}_2)_z-$  where  $\text{X}_e$  is

5  $\text{C}(\text{R}_{8m})(\text{R}_{8l})$ ;









$\text{R}_{8f}$  is alkyl or aryl.

13. The compound as defined in Claim 1 having the structure































5







5









5







Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral



Chiral






Chiral




Chiral




















5




































































= X<sub>1</sub>-R

X<sub>1</sub>-R =



















 $= X_1-R$  $X_1-R =$ 





$X_1-R =$








































X<sub>1</sub>-R =



X<sub>1</sub>-CH<sub>3</sub>,







Chemical structures of various substituted compounds, including aromatic and aliphatic derivatives, with substituents like  $X_1$ ,  $H_3C$ ,  $CH_3$ ,  $OCH_3$ ,  $F$ , and  $CF_3$ .

Row 1:

- $X_1-CH_2-C_6H_5$
- $X_1-C_6H_{11}$
- $X_1-C_6H_4-CH_3$

Row 2:

- $X_1-C_6H_3(CH_3)_2$
- $X_1-C_6H_5$
- $X_1-CH_2-C(CH_3)_2-CH_3$

Row 3:

- $X_1-C_6H_4-OCH_3$
- $X_1-C_6H_4-OCH_3$
- $X_1-C_6H_5$

Row 4:

- $X_1-CH_2-CH_2-CF_3$
- $X_1-C_6H_4-OCH_3$
- $X_1-C_6H_4-F$





























Row 5:













- $X_1-CH_2-CH_2-CH_3$
- $X_1-C_6H_4-CF_3$
- $X_1-C_6H_4-CF_3$

Row 6:

- $X_1-C_6H_4-CH_3$
- $X_1-C_6H_4-CO-C_6H_5$
- $X_1-CH_2-CH=CH-CH_3$























































**X<sub>1</sub> - R**





= X<sub>1</sub>-R



X<sub>1</sub>-R











































 $X_1-R$ 


5

 $X_1-R$ 


X1CCCCNC(=O)OCCO




## Structure




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral



Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral




Chiral



Chiral




Chiral



or a pharmaceutically acceptable salt thereof.


14. The compound as defined in Claim 1 having the structure

5




10








Chiral




Chiral




Chiral




Chiral





Chiral




Chiral




Chiral




Chiral




Chiral



Chiral






Chiral




Chiral

5







Chiral




Chiral




Chiral




Chiral




Chiral



Chiral



Chiral




Chiral


5

or a pharmaceutically acceptable salt thereof.

15. The compound as defined in Claim 1 having the structure




Chiral



Chiral



Chiral




Chiral



or a pharmaceutically acceptable salt thereof.

16. The compound as defined in Claim 15 having the structure

5




10

, or a pharmaceutically acceptable salt thereof.

17. The compound as defined in Claim 15 having the structure

15



or

or a pharmaceutically acceptable salt thereof.

18. A pharmaceutical composition comprising a compound as defined in Claim 1 and a pharmaceutically acceptable carrier therefor.

20

19. A method for increasing levels of endogenous growth hormone, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

5

20. A method for treating obesity, osteoporosis, renal disease, cardiac myopathy, cachexia, HIV wasting syndrome, long term critical illness, sarcopenia, and/or stimulating wound healing and/or the immune system, or increasing muscle mass and/or muscle strength, or maintenance of muscle strength and function in the elderly, or reversal or prevention of frailty in the elderly, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

10

15

21. A method for treating Syndrome X, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

20

22. A method for prophylaxis and/or treatment of diabetes and/or increasing lean body mass, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

25

23. A method for preventing or treating osteoporosis, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

30

24. A method for treating osteoporosis, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1 in combination with parathyroid hormone or a bisphosphonate.

35

25. A method for treating Syndrome X, cachexia, HIV wasting syndrome, long term critical illness, or sarcopenia, or for increasing muscle mass and/or muscle strength, or for maintenance of muscle strength and function in the elderly, or for reversal or prevention of frailty in the elderly, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1 in combination with estrogen, testosterone, a selective estrogen receptor modulator, or a selective androgen receptor modulator.